

■Chapter 1. Definition

NISHIYAMA Toshio

Professor Emeritus, Tokyo University of Agriculture and Technology

- 1) 林利彦. コラーゲン線維形成. 日本生物物理学会 編. 生命科学の基礎3 形態形成. 東京大学出版; 1977. p. 83-124.
- 2) 服部俊治. 動物由来線維分子コラーゲンの性質と応用. *Sen'i Gakkaishi (繊維と工業)*. 2009; 65: 453-461.
- 3) 横山隆亮, 西谷和彦. 植物細胞壁の構造と機能の多様性. *植物科学最前線*. 2014; 5: 45-52.
- 4) King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, et al. The genome of the choanoflagellate *monosiga brevicollis* and the origin of metazoans. *Nature* 2008; 451: 783-8.
- 5) Fidler AL, Darris CE, Chetyrkin SV, Pedchenko VK, Boudko SP, Brown KL, et al. Collagen IV and basement membrane at the evolutionary dawn of metazoan tissues. *Elife* 2017; 6: e24176.
- 6) Ryan JF, Pang K, Schnitzler CE, Nguyen AD, Moreland RT, Simmons DK, et al. The genome of the ctenophore *mnemiopsis leidyi* and its implications for cell type evolution. *Science* 2013; 342: 1242592.
- 7) Grau-Bove X, Torruella G, Donachie S, Suga H, Leonard G, Richards TA, et al. Dynamics of genomic innovation in the unicellular ancestry of animals. *Elife* 2017; 6: e26036.
- 8) Fidler AL, Boudko SP, Rokas A, Hudson BG. The triple helix of collagens – an ancient protein structure that enabled animal multicellularity and tissue evolution. *J Cell Sci.* 2018; 131: jcs203950.
- 9) Aouacheria A, Geourjon C, Aghajari N, Navratil V, Deléage G, Lethias C, et al. Insights into early extracellular matrix evolution: spongin short chain collagen-related proteins are homologous to basement membrane type IV collagens and form a novel family widely distributed in invertebrates. *Mol Biol Evol.* 2006; 23: 2288-2302.
- 10) 松村外志張. コラーゲンの比較生化学. 野田, 永井, 藤本 編. コラーゲン. 南江堂; 1975. p. 169-206.
- 11) 木村茂. 水産動物のコラーゲン. *化学と生物*. 1987; 25: 311-7.
- 12) 吉里勝利. コラーゲンの比較生化学. *比較生理生化学*. 1991; 8: 72-80.
- 13) 安達栄治郎, 澤田元. 細胞外マトリックスの主要線維成分. 小出, 林 編. 細胞外マトリックス –基礎と臨床–. 愛智出版; 2000. p. 2-40.
- 14) 二宮善文, 吉岡秀克. コラーゲンスーパーファミリーの遺伝子群. 小出, 林 編. 細胞外マトリックス –基礎と臨床–. 愛智出版; 2000. p. 53-93.
- 15) 水野一乗, 林利彦. 1章 コラーゲン 基礎知識. 畑, 服部, 新井 編. 細胞外マトリックス研究法 [1] 第1部. 生化学的研究法. コラーゲン技術研修会; 1998. P. 5-13.
- 16) 林利彦, 水野一乗, 中里浩一, 羽関典子, 西山敏夫. コラーゲンスーパーファミリータンパク質. 小出, 林 編. 細胞外マトリックス –基礎と臨床–. 愛智出版; 2000. p. 94-138.

- 17) 西山敏夫. 表面と表面の相互作用が⁶生体機能を制御する. 表面. 2010; 48: 285-287.
- 18) 富永直樹, 西山敏夫, 林利彦. 皮膚とコラーゲン. *Sen'i Gakkaishi (繊維と工業)*. 1986; 42: 237-242.
- 19) 西田紀貴, 嶋田一夫. コラーゲン結合タンパク質を介した生命プロセスの活性化機構. *生化学*. 2008; 80: 483-492.
- 20) Kadler KE, Baldock C, Bella J, Boot-Handford RP. Collagens at a glance. *J Cell Sci*. 2007; 120: 1955-8.
- 21) Gordon MK, Hahn RA. Collagens. *Cell Tissue Res*. 2010; 339: 247-257.
- 22) Ricard-Blum, S. The collagen family. *Cold Spring Harb Perspect Biol*. 2011; 3: a004978.
- 23) 奥山健二, 川口辰也. コラーゲンの分子構造・高次構造. *高分子論文集*. 2010; 67: 229-247.
- 24) 奥山健二. コラーゲンの構造. *日本結晶学会誌*. 2012; 54:263-269.
- 25) Söderhäll C, Marenholz I, Kerscher T, Rüschenhoff F, Esparza-Gordillo J, Worm M, et al. Variants in a novel epidermal collagen gene (COL29A1) are associated with atopic dermatitis. *PLoS Biol*. 2007; 5: e242.
- 26) Gara SK, Grumati P, Urciuolo A, Bonaldo P, Kobbe B, Koch M, et al. Three novel collagen VI chains with high homology to the alpha3 chain. *J Biol Chem*. 2008; 283: 10658-10670.
- 27) Borza DB, Bondar O, Ninomiya Y, Sado Y, Naito I, Todd P, et al. The NC1 domain of collagen IV encodes a novel network composed of the alpha 1, alpha 2, alpha 5, and alpha 6 chains in smooth muscle basement membranes. *J Biol Chem*. 2001; 276: 28532-28540.

■Chapter 2. Collagen Family Proteins

MIZUNO Kazunori

Nippi Research Institute of Biomatrix

- 1) Bächinger HP, Mizuno K, Vranka J, and Boudko SP. Collagen formation and structure. In *Comprehensive natural products II Chemistry and Biology* Mander L, Lui H-W, editors, Elsevier; Oxford; 2010. p. 469–530.
- 2) Kadler KE, Baldock C, Bella J, and Boot-Handford RP. Collagens at a glance. *J Cell Sci.* 2007; 120: 1955–8.
- 3) Kielty CM, Hopkinson I, and Grant ME. The collagen family: structure, assembly, and organization in the extracellular matrix. In *Connective Tissue and Its Heritable Disorders* Royce PM, Steinmann B, editors, 2nd Ed., Wiley-Liss; New York, USA; 2002. p. 159–221.
- 4) Ricard-Blum S. The collagen family. *Cold Spring Harb Perspect Biol.* 2011; 3: a004978.
- 5) Holmes DF, Lu Y, Starborg T, and Kadler KE. Collagen Fibril Assembly and Function. *Curr Top Dev Biol.* 2018; 130: 107–42.
- 6) Kadler KE. Fell Muir Lecture: Collagen fibril formation in vitro and in vivo. *Int J Exp Pathol.* 2017; 98: 4–16.
- 7) Kadler KE, Hill A, and Canty-Laird EG. Collagen fibrillogenesis: fibronectin, integrins, and minor collagens as organizers and nucleators. *Curr Opin Cell Biol.* 2008; 20: 495–501.
- 8) Svensson RB, Herchenhan A, Starborg T, Larsen M, Kadler KE, Qvortrup K, et al. Evidence of structurally continuous collagen fibrils in tendons. *Acta Biomater.* 2017; 50: 293–301.
- 9) Byers PH, and Cole WG. Osteogenesis imperfecta. In *Connective Tissue and Its Heritable Disorders* Royce PM, Steinmann B, editors, 2nd Ed., Wiley-Liss, New York, USA; 2002. p. 385–430.
- 10) Makareeva E, Mertz EL, Kuznetsova NV, Sutter MB, DeRidder AM, Cabral WA, et al. Structural heterogeneity of type I collagen triple helix and its role in osteogenesis imperfecta. *J Biol Chem.* 2008; 283: 4787–98.
- 11) Han S, Makareeva E, Kuznetsova NV, DeRidder AM, Sutter MB, Losert W, et al. Molecular mechanism of type I collagen homotrimer resistance to mammalian collagenases. *J Biol Chem.* 2010; 285: 22276–81.
- 12) Makareeva E, Han S, Vera JC, Sackett DL, Holmbeck K, Phillips CL, et al. Carcinomas contain a matrix metalloproteinase-resistant isoform of type I collagen exerting selective support to invasion. *Cancer Res.* 2010; 70: 4366–74.
- 13) Miller EJ, and Matukas VJ. Chick cartilage collagen: a new type of alpha 1 chain not present in bone or skin of the species. *Proc Natl Acad Sci U S A.* 1969; 64: 1264–8.
- 14) Keene DR, Sakai LY, Bächinger HP, and Burgeson RE. Type III collagen can be present on banded collagen fibrils regardless of fibril diameter. *J Cell Biol.* 1987; 105: 2393–402.

- 15) Liu X, Wu H, Byrne M, Krane S, and Jaenisch R. Type III collagen is crucial for collagen I fibrillogenesis and for normal cardiovascular development. *Proc Natl Acad Sci U S A*. 1997; 94: 1852-6.
- 16) Eyre DR, Weis MA, and Wu JJ. Articular cartilage collagen: an irreplaceable framework? *Eur Cell Mater*. 2006; 12: 57-63.
- 17) Brisson BK, Mauldin EA, Lei W, Vogel LK, Power AM, Lo A, et al. Type III Collagen Directs Stromal Organization and Limits Metastasis in a Murine Model of Breast Cancer. *Am J Pathol*. 2015; 185: 1471-86.
- 18) Fessler JH, and Fessler LI. Type V collagen. In *Structure and function of collagen types* Mayne R, Burgeson RE, editors, Academic Press, Orlando, FL, USA; 1987. p. 81-103.
- 19) Fichard A, Kleman JP, and Ruggiero F. Another look at collagen V and XI molecules. *Matrix Biol*. 1995; 14: 515-31.
- 20) Adachi E, Hopkinson I, and Hayashi T. Basement-membrane stromal relationships: interactions between collagen fibrils and the lamina densa. *Int Rev Cytol*. 1997; 173: 73-156.
- 21) Niyibizi C, Fietzek PP, and van der Rest M. Human placenta type V collagens. Evidence for the existence of an alpha 1(V) alpha 2(V) alpha 3(V) collagen molecule. *J Biol Chem*. 1984; 259: 14170-4.
- 22) Adachi E, and Hayashi T. In vitro formation of fine fibrils with a D-periodic banding pattern from type V collagen. *Coll Relat Res*. 1985; 5: 225-32.
- 23) Adachi E, and Hayashi T. In vitro formation of hybrid fibrils of type V collagen and type I collagen. Limited growth of type I collagen into thick fibrils by type V collagen. *Connect Tissue Res*. 1986; 14: 257-66.
- 24) Huang G, Ge G, Izzi V, and Greenspan DS. alpha3 Chains of type V collagen regulate breast tumour growth via glypican-1. *Nat Commun*. 2017; 8: 14351.
- 25) Huang G, Ge G, Wang D, Gopalakrishnan B, Butz DH, Colman RJ, et al. alpha3(V) collagen is critical for glucose homeostasis in mice due to effects in pancreatic islets and peripheral tissues. *J Clin Invest*. 2011; 121: 769-83.
- 26) Eyre D, and Wu JJ. Type XI collagen. In *Structure and function of collagen types* Mayne R, Burgeson RE, editors, Academic Press, Orlando, FL, USA; 1987. p.
- 27) Morris NP, and Bächinger HP. Type XI collagen is a heterotrimer with the composition (1 alpha, 2 alpha, 3 alpha) retaining non-triple-helical domains. *J Biol Chem*. 1987; 262: 11345-50.
- 28) Burgeson RE, and Hollister DW. Collagen heterogeneity in human cartilage: identification of several new collagen chains. *Biochem Biophys Res Commun*. 1979; 87: 1124-31.
- 29) Thom JR, and Morris NP. Biosynthesis and proteolytic processing of type XI collagen in embryonic chick sterna. *J Biol Chem*. 1991; 266: 7262-9.
- 30) Koch M, Laub F, Zhou P, Hahn RA, Tanaka S, Burgeson RE, et al. Collagen XXIV, a vertebrate

fibrillar collagen with structural features of invertebrate collagens: selective expression in developing cornea and bone. *J Biol Chem.* 2003; 278: 43236–44.

- 31) Hjorten R, Hansen U, Underwood RA, Telfer HE, Fernandes RJ, Krakow D, et al. Type XXVII collagen at the transition of cartilage to bone during skeletogenesis. *Bone.* 2007; 41: 535–42.
- 32) Boutaud A, Borza DB, Bondar O, Gunwar S, Netzer KO, Singh N, et al. Type IV collagen of the glomerular basement membrane. Evidence that the chain specificity of network assembly is encoded by the noncollagenous NC1 domains. *J Biol Chem.* 2000; 275: 30716–24.
- 33) Sado Y, Kagawa M, Naito I, Ueki Y, Seki T, Momota R, et al. Organization and expression of basement membrane collagen IV genes and their roles in human disorders. *J Biochem.* 1998; 123: 767–76.
- 34) Saito K, Naito I, Seki T, Oohashi T, Kimura E, Momota R, et al. Differential expression of mouse alpha5(IV) and alpha6(IV) collagen genes in epithelial basement membranes. *J Biochem.* 2000; 128: 427–34.
- 35) Hasegawa H, Naito I, Nakano K, Momota R, Nishida K, Taguchi T, et al. The distributions of type IV collagen alpha chains in basement membranes of human epidermis and skin appendages. *Arch Histol Cytol.* 2007; 70: 255–65.
- 36) Nakano K, Naito I, Momota R, Sado Y, Hasegawa H, Ninomiya Y, et al. The distribution of type IV collagen alpha chains in the mouse ovary and its correlation with follicular development. *Arch Histol Cytol.* 2007; 70: 243–53.
- 37) Nakano S, Iyama K, Ogawa M, Yoshioka H, Sado Y, Oohashi T, et al. Differential tissular expression and localization of type IV collagen alpha1(IV), alpha2(IV), alpha5(IV), and alpha6(IV) chains and their mRNA in normal breast and in benign and malignant breast tumors. *Lab Invest.* 1999; 79: 281–92.
- 38) Pedchenko V, Bauer R, Pokidysheva EN, Al-Shaer A, Forde NR, Fidler AL, et al. A chloride ring is an ancient evolutionary innovation mediating the assembly of the collagen IV scaffold of basement membranes. *J Biol Chem.* 2019; 294: 7968–81.
- 39) Robertson WE, Rose KL, Hudson BG, and Vanacore RM. Supramolecular organization of the alpha121–alpha565 collagen IV network. *J Biol Chem.* 2014; 289: 25601–10.
- 40) Adachi E, Takeda Y, Nakazato K, Muraoka M, Iwata M, Sasaki T, et al. Isolated collagen IV retains the potential to form an 18-nm sided polygonal meshwork of the lamina densa. *J Electron Microsc (Tokyo).* 1997; 46: 233–41.
- 41) Vanacore RM, Friedman DB, Ham A–JL, Sundaramoorthy M, and Hudson BG. Identification of S-hydroxylysyl-methionine as the covalent cross-link of the noncollagenous (NC1) hexamer of the alpha1alpha1alpha2 collagen IV network: a role for the post-translational modification of lysine 211 to hydroxylysine 211 in hexamer assembly. *J Biol Chem.* 2005; 280: 29300–10.
- 42) Fidler AL, Darris CE, Chetyrkin SV, Pedchenko VK, Boudko SP, Brown KL, et al. Collagen IV and

basement membrane at the evolutionary dawn of metazoan tissues. *Elife*. 2017; 6:

- 43) Matsuura-Hachiya Y, Arai KY, Muraguchi T, Sasaki T, and Nishiyama T. Type IV collagen aggregates promote keratinocyte proliferation and formation of epidermal layer in human skin equivalents. *Exp Dermatol*. 2018; 27: 443–8.
- 44) Gordon MK, Gerecke DR, Dublet B, van der Rest M, and Olsen BR. Type XII collagen. A large multidomain molecule with partial homology to type IX collagen. *J Biol Chem*. 1989; 264: 19772–8.
- 45) Dublet B, and van der Rest M. Type XIV collagen, a new homotrimeric molecule extracted from fetal bovine skin and tendon, with a triple helical disulfide-bonded domain homologous to type IX and type XII collagens. *J Biol Chem*. 1991; 266: 6853–8.
- 46) Kassner A, Tiedemann K, Notbohm H, Ludwig T, Morgelin M, Reinhardt DP, et al. Molecular structure and interaction of recombinant human type XVI collagen. *J Mol Biol*. 2004; 339: 835–53.
- 47) Boudko SP, and Bachinger HP. The NC2 domain of type IX collagen determines the chain register of the triple helix. *J Biol Chem*. 2012; 287: 44536–45.
- 48) Boudko SP, Zientek KD, Vance J, Hacker JL, Engel J, and Bachinger HP. The NC2 domain of collagen IX provides chain selection and heterotrimerization. *J Biol Chem*. 2010; 285: 23721–31.
- 49) Chiquet M, Birk DE, Bonnemann CG, and Koch M. Collagen XII: Protecting bone and muscle integrity by organizing collagen fibrils. *Int J Biochem Cell Biol*. 2014; 53: 51–4.
- 50) Nishiyama T, McDonough AM, Bruns RR, and Burgeson RE. Type XII and XIV collagens mediate interactions between banded collagen fibers in vitro and may modulate extracellular matrix deformability. *J Biol Chem*. 1994; 269: 28193–9.
- 51) Akutsu N, Milbury CM, Burgeson RE, and Nishiyama T. Effect of type XII or XIV collagen NC-3 domain on the human dermal fibroblast migration into reconstituted collagen gel. *Exp Dermatol*. 1999; 8: 17–21.
- 52) Kuo HJ, Maslen GL, Keene DR, and Glanville RW. Type VI collagen anchors endothelial basement membranes by interacting with type IV collagen. *J Biol Chem*. 1997; 272: 26522–9.
- 53) Timple R, and Engel J. Type VI collagen. In *Structure and function of collagen types* Mayne R, Burgeson RE, editors, Academic Press, Orlando, FL, USA; 1987. p.
- 54) Keene DR, Sakai LY, Lunstrum GP, Morris NP, and Burgeson RE. Type VII collagen forms an extended network of anchoring fibrils. *J Cell Biol*. 1987; 104: 611–21.
- 55) Fitzgerald J, Rich C, Zhou FH, and Hansen U. Three novel collagen VI chains, alpha 4(VI), alpha 5(VI) and alpha 6(VI). *J Biol Chem*. 2008;
- 56) Gara SK, Grumati P, Urciuolo A, Bonaldo P, Kobbe B, Koch M, et al. Three novel collagen VI chains with high homology to the alpha3 chain. *J Biol Chem*. 2008; 283: 10658–70.

- 57) Wiberg C, Heinegard D, Wenglen C, Timpl R, and Morgelin M. Biglycan organizes collagen VI into hexagonal-like networks resembling tissue structures. *J Biol Chem.* 2002; 277: 49120–6.
- 58) Godwin ARF, Starborg T, Sherratt MJ, Roseman AM, and Baldock C. Defining the hierarchical organisation of collagen VI microfibrils at nanometre to micrometre length scales. *Acta Biomater.* 2017; 52: 21–32.
- 59) Lunstrum GP, Kuo HJ, Rosenbaum LM, Keene DR, Glanville RW, Sakai LY, et al. Anchoring fibrils contain the carboxyl-terminal globular domain of type VII procollagen, but lack the amino-terminal globular domain. *J Biol Chem.* 1987; 262: 13706–12.
- 60) Lunstrum GP, Sakai LY, Keene DR, Morris NP, and Burgeson RE. Large complex globular domains of type VII procollagen contribute to the structure of anchoring fibrils. *J Biol Chem.* 1986; 261: 9042–8.
- 61) Sakai LY, Keene DR, Morris NP, and Burgeson RE. Type VII collagen is a major structural component of anchoring fibrils. *J Cell Biol.* 1986; 103: 1577–86.
- 62) van den Akker PC, Jonkman MF, Rengaw T, Bruckner-Tuderman L, Has C, Bauer JW, et al. The international dystrophic epidermolysis bullosa patient registry: an online database of dystrophic epidermolysis bullosa patients and their COL7A1 mutations. *Hum Mutat.* 2011; 32: 1100–7.
- 63) Mann K, Jander R, Korsching E, Kühn K, and Rauterberg J. The primary structure of a triple-helical domain of collagen type VIII from bovine Descemet's membrane. *FEBS Lett.* 1990; 273: 168–72.
- 64) Muragaki Y, Jacenko O, Apte S, Mattei MG, Ninomiya Y, and Olsen BR. The alpha 2(VIII) collagen gene. A novel member of the short chain collagen family located on the human chromosome 1. *J Biol Chem.* 1991; 266: 7721–7.
- 65) Muragaki Y, Shiota C, Inoue M, Ooshima A, Olsen BR, and Ninomiya Y. alpha 1(VIII)-collagen gene transcripts encode a short-chain collagen polypeptide and are expressed by various epithelial, endothelial and mesenchymal cells in newborn mouse tissues. *Eur J Biochem.* 1992; 207: 895–902.
- 66) Myers JC, Amenta PS, Dion AS, Sciancalepore JP, Nagaswami C, Weisel JW, et al. The molecular structure of human tissue type XV presents a unique conformation among the collagens. *Biochem J.* 2007; 404: 535–44.
- 67) Myers JC, Dion AS, Abraham V, and Amenta PS. Type XV collagen exhibits a widespread distribution in human tissues but a distinct localization in basement membrane zones. *Cell Tissue Res.* 1996; 286: 493–505.
- 68) Amenta PS, Scivoletti NA, Newman MD, Sciancalepore JP, Li D, and Myers JC. Proteoglycan-collagen XV in human tissues is seen linking banded collagen fibers subjacent to the basement membrane. *J Histochem Cytochem.* 2005; 53: 165–76.

- 69) Sasaki T, Larsson H, Tisi D, Claesson-Welsh L, Hohenester E, and Timpl R. Endostatins derived from collagens XV and XVIII differ in structural and binding properties, tissue distribution and anti-angiogenic activity. *J Mol Biol.* 2000; 301: 1179–90.
- 70) Sasaki T, Fukai N, Mann K, Göhring W, Olsen BR, and Timpl R. Structure, function and tissue forms of the C-terminal globular domain of collagen XVIII containing the angiogenesis inhibitor endostatin. *EMBO J.* 1998; 17: 4249–56.
- 71) Heljasvaara R, Aikio M, Ruotsalainen H, and Pihlajaniemi T. Collagen XVIII in tissue homeostasis and dysregulation – Lessons learned from model organisms and human patients. *Matrix Biol.* 2017; 57–58: 55–75.
- 72) Oh SP, Warman ML, Seldin MF, Cheng SD, Knoll JH, Timmons S, et al. Cloning of cDNA and genomic DNA encoding human type XVIII collagen and localization of the alpha 1(XVIII) collagen gene to mouse chromosome 10 and human chromosome 21. *Genomics.* 1994; 19: 494–9.
- 73) Boudko SP, Sasaki T, Engel J, Lerch TF, Nix J, Chapman MS, et al. Crystal structure of human collagen XVIII trimerization domain: A novel collagen trimerization fold. *J Mol Biol.* 2009; 392: 787–802.
- 74) Wirz JA, Boudko SP, Lerch TF, Chapman MS, and Bachinger HP. Crystal structure of the human collagen XV trimerization domain: a potent trimerizing unit common to multiplexin collagens. *Matrix Biol.* 2011; 30: 9–15.
- 75) Li D, Clark CC, and Myers JC. Basement membrane zone type XV collagen is a disulfide-bonded chondroitin sulfate proteoglycan in human tissues and cultured cells. *J Biol Chem.* 2000; 275: 22339–47.
- 76) Sasaki T, Hohenester E, and Timpl R. Structure and function of collagen-derived endostatin inhibitors of angiogenesis. *IUBMB Life.* 2002; 53: 77–84.
- 77) Marneros AG, and Olsen BR. Physiological role of collagen XVIII and endostatin. *FASEB J.* 2005; 19: 716–28.
- 78) Franzke CW, Bruckner P, and Bruckner-Tuderman L. Collagenous transmembrane proteins: recent insights into biology and pathology. *J Biol Chem.* 2005; 280: 4005–8.
- 79) Peltonen S, Rehn M, and Pihlajaniemi T. Alternative splicing of mouse alpha1(XIII) collagen RNAs results in at least 17 different transcripts, predicting alpha1(XIII) collagen chains with length varying between 651 and 710 amino acid residues. *DNA Cell Biol.* 1997; 16: 227–34.
- 80) Hägg P, Rehn M, Huhtala P, Väisänen T, Tamminen M, and Pihlajaniemi T. Type XIII collagen is identified as a plasma membrane protein. *J Biol Chem.* 1998; 273: 15590–7.
- 81) Sandberg-Lall M, Hägg PO, Wahlström I, and Pihlajaniemi T. Type XIII collagen is widely expressed in the adult and developing human eye and accentuated in the ciliary muscle, the optic nerve and the neural retina. *Exp Eye Res.* 2000; 70: 401–10.
- 82) Matsumura H, Mohri Y, Binh NT, Morinaga H, Fukuda M, Ito M, et al. Hair follicle aging is

- driven by transepidermal elimination of stem cells via COL17A1 proteolysis. *Science*. 2016; 351: aad4395.
- 83) Liu N, Matsumura H, Kato T, Ichinose S, Takada A, Namiki T, et al. Stem cell competition orchestrates skin homeostasis and ageing. *Nature*. 2019; 568: 344–50.
 - 84) Hashimoto T, Wakabayashi T, Watanabe A, Kowa H, Hosoda R, Nakamura A, et al. CLAC: a novel Alzheimer amyloid plaque component derived from a transmembrane precursor, CLAC-P/collagen type XXV. *EMBO J*. 2002; 21: 1524–34.
 - 85) Munezane H, Oizumi H, Wakabayashi T, Nishio S, Hirasawa T, Sato T, et al. Roles of Collagen XXV and Its Putative Receptors PTPsigma/delta in Intramuscular Motor Innervation and Congenital Cranial Dysinnervation Disorder. *Cell Rep*. 2019; 29: 4362–76 e6.
 - 86) Goncalves TJM, Boutillon F, Lefebvre S, Goffin V, Iwatsubo T, Wakabayashi T, et al. Collagen XXV promotes myoblast fusion during myogenic differentiation and muscle formation. *Sci Rep*. 2019; 9: 5878.
 - 87) Tanaka T, Wakabayashi T, Oizumi H, Nishio S, Sato T, Harada A, et al. CLAC-P/collagen type XXV is required for the intramuscular innervation of motoneurons during neuromuscular development. *J Neurosci*. 2014; 34: 1370–9.
 - 88) Sato K, Yomogida K, Wada T, Yorihuzi T, Nishimune Y, Hosokawa N, et al. Type XXVI collagen, a new member of the collagen family, is specifically expressed in the testis and ovary. *J Biol Chem*. 2002; 277: 37678–84.
 - 89) Veit G, Kobbe B, Keene DR, Paulsson M, Koch M, and Wagener R. Collagen XXVIII, a novel von Willebrand factor A domain-containing protein with many imperfections in the collagenous domain. *J Biol Chem*. 2006; 281: 3494–504.
 - 90) Turner MW. The role of mannose-binding lectin in health and disease. *Mol Immunol*. 2003; 40: 423–9.
 - 91) Vieira F, Kung JW, and Bhatti F. Structure, genetics and function of the pulmonary associated surfactant proteins A and D: The extra-pulmonary role of these C type lectins. *Ann Anat*. 2017; 211: 184–201.
 - 92) Hoppe HJ, and Reid KB. Collectins—soluble proteins containing collagenous regions and lectin domains—and their roles in innate immunity. *Protein Sci*. 1994; 3: 1143–58.
 - 93) Casals C, Garcia-Fojeda B, and Minutti CM. Soluble defense collagens: Sweeping up immune threats. *Mol Immunol*. 2019; 112: 291–304.
 - 94) Takamatsu N, Ohba K, Kondo J, Kondo N, and Shiba T. Hibernation-associated gene regulation of plasma proteins with a collagen-like domain in mammalian hibernators. *Mol Cell Biol*. 1993; 13: 1516–21.
 - 95) Ruotsalainen H, Risteli M, Wang C, Wang Y, Karppinen M, Bergmann U, et al. The activities of lysyl hydroxylase 3 (LH3) regulate the amount and oligomerization status of adiponectin. *PLoS*

One. 2012; 7: e50045.

- 96) Doi T, Higashino K, Kurihara Y, Wada Y, Miyazaki T, Nakamura H, et al. Charged collagen structure mediates the recognition of negatively charged macromolecules by macrophage scavenger receptors. *J Biol Chem.* 1993; 268: 2126–33.

■Chapter 3. The Structure and Physical Properties of Collagen

NOMURA Yoshihiro

Scleroprotein and Leather Research Institute, Faculty of Agriculture, Tokyo University of
Agriculture and Technology

GOH K. L.

Newcastle University in Singapore, Singapore

Newcastle University, Newcastle Upon Tyne, UK

- 1) GN Ramachandran and G Kartha, Structure of collagen. *Nature* 176, 593–595 (1955)
- 2) A Rich and H Crick The molecular structure of collagen. *J Mol Biol*, 3, 483–506 (1961)
- 3) KL Goh, As tough as leather: Macro to nano scale perspectives of collagen stability. IULTCS Congress. 1–17 (2019).
- 4) L Arseni, A Lombardi, D Orioli. From structure to phenotype: Impact of collagen alterations on human health. *Molecular Sci*, 19, 1407 (2018)
- 5) 奥山隆、川口辰也、コラーゲンの分子構造・高次構造 高分子論文集, 67(4). 229–247 (2010)
- 6) MD Shoulders, RT Raines. Collagen structure and stability. *Annu Rev Biochem*, 78: 929–958 (2009)
- 7) J Emsley, CG Knight, RW Farndale, MJ Barnes, RC Liddington. Structural basis of collagen recognition by integrin $\alpha 2\beta 1$. *Cell*, 101: 47–56 (2000)
- 8) SM Sweeney, CA Guy, GB Fields, JD San Antonio. Defining the domains of type I collagen involved in heparin-binding and endothelial tube formation. *Proc. Natl. Acad. Sci. USA*, 95: 7275–7280 (1998).
- 9) MJ Buehler. Nature designs tough collagen: explaining the nanostructure of collagen fibrils. *Proc. Natl. Acad. Sci. USA*, 103: 12285–12290 (2006).
- 10) J Howard. *Mechanics of Motor Proteins and the Cytoskeleton*. Sunderland, MA: Sinauer; 2001
- 11) PJ in 't Veld, MJ Stevens. Simulation of the mechanical strength of a single collagen molecule. *Biophys. J*, 95: 33–39 (2008)
- 12) JAJ van der Rijt, KO van der Werf, ML Bennink, PJ Dijkstra, J Feijen. Micromechanical testing of individual collagen fibrils. *Macromol. Biosci*, 6: 697–702 (2006)
- 13) MPE Wenger, L Bozec, M Horton, P Mesquida. Mechanical properties of collagen fibrils. *Biophys. J*, 93: 1255–1263 (2007)
- 14) Zhang Y, Wang L. Recent research progress on leather fat liquoring agents. *Polymer-Plastics Technology and Engineering*, 48: 285–291 (2009)
- 15) NT Wright, JD Humphrey: Denaturation of collagen via heating: an irreversible rate process. *Annu Rev Biomed Eng*, 4: 109–128 (2002)
- 16) YL Yeo, KL Goh, L Kin, HJ Wang, A Listrat, D Bechet. Structure–property relationship of burn

- collagen reinforcing musculo-skeletal tissues. *Key Eng Mater*, 478:87–92 (2011)
- 17) KS Weadock, EJ Miller, LD Bellincampi, JP Zawadsky, MG Dunn. Physical crosslinking of collagen fibers – comparison of ultraviolet irradiation and dehydrothermal treatment. *Journal of Biomedical Materials Research*, 29:1373–1379 (1995)
 - 18) KL Goh, DF Holmes, HY Lu, S Richardson, KE Kadler, PP Purslow, TJ Wess. Ageing changes in the tensile properties of tendons: influence of collagen fibril volume. *J Biomech Eng*, 130:021011 (2008)
 - 19) B Gogly, G Godeau, S Gilbert, JM Legrand, C Kut, B Pellat, M Goldberg. Morphometric analysis of collagen and elastic fibers in normal skin and gingiva in relation to age. *Clinical Oral Investigation*, 1:147–152 (1997)
 - 20) HC Wells, KH Sizeland, HR Kaye, N Kirby, A Hawley, ST Mudie, RG Haverkamp. Poisson's ratio of collagen fibrils measured by small angle X-ray scattering of strained bovine pericardium. *J Appl Phys*, 117: 044701 (2015)
 - 21) J Liao, I Vesely. A structural basis for the size-related mechanical properties of mitral valve chordae tendineae. *Journal of Biomechanics*, 36(8): 1125–1133 (2003)
 - 22) C Morin, C Hellmich, P Henits. Fibrillar structure and elasticity of hydrating collagen: a quantitative multiscale approach. *J Theor Biol*, 317: 384–393 (2013)
 - 23) HC Wells, KH Sizeland, N Kirby, A Hawley, S Mudie, RG Haverkamp. Acellular dermal matrix collagen responds to strain by intermolecular spacing contraction with fibril extension and rearrangement. *J Mech Behav Biomed Mater*, 79: 1–8 (2018)
 - 24) HC Wells, G Holmes, RG Haverkamp. Looseness in bovine leather: microstructural characterization. *J Sci Food Agric*, 96(8): 2731–2736 (2016)
 - 25) SJ Kelly, HC Wells, KH Sizeland, N Kirby, RL Edmonds, T Ryan, A Hawley, S Mudie, RG Haverkamp. Artificially modified collagen fibril orientation affects leather tear strength. *J Sci Food Agric*, 98(9): 3524–3531 (2018)

■Chapter 4. Function in vivo

MIYATA Shinji

Scleroprotein and Leather Research Institute, Faculty of Agriculture, Tokyo University of Agriculture and Technology

- 1) Löhler J, Timpl R, Jaenisch R. Embryonic lethal mutation in mouse collagen I gene causes rupture of blood vessels and is associated with erythropoietic and mesenchymal cell death. *Cell*. 1984; 38: 597-607.
- 2) Bonadio J, Saunders TL, Tsai E, Goldstein SA, Morris-Wiman J, Brinkley L, et al. Transgenic mouse model of the mild dominant form of osteogenesis imperfecta. *Proc Natl Acad Sci U S A*. 1990; 87: 7145-7149.
- 3) Gajko-Galicka A. Mutations in type I collagen genes resulting in osteogenesis imperfecta in humans. *Acta Biochim Pol*. 2002; 49: 433-441.
- 4) Spotila LD, Constantinou CD, Sereda L, Ganguly A, Riggs BL, Prockop DJ. Mutation in a gene for type I procollagen (COL1A2) in a woman with postmenopausal osteoporosis: evidence for phenotypic and genotypic overlap with mild osteogenesis imperfecta. *Proc Natl Acad Sci U S A*. 1991; 88: 5423-5427.
- 5) Huang G, Ge G, Wang D, Gopalakrishnan B, Butz DH, Colman RJ, et al. $\alpha 3(V)$ Collagen is critical for glucose homeostasis in mice due to effects in pancreatic islets and peripheral tissues. *J Clin Invest*. 2011; 121: 769-783.
- 6) Chiquet M, Birk DE, Bönemann CG, Koch M. Collagen XII: Protecting bone and muscle integrity by organizing collagen fibrils. *Int J Biochem Cell Biol*. 2014; 53: 51-54.
- 7) Izu Y, Sun M, Zwolanek D, Veit G, Williams V, Cha B, et al. Type XII collagen regulates osteoblast polarity and communication during bone formation. *J Cell Biol*. 2011; 193: 1115-1130.
- 8) Ansorge HL, Meng X, Zhang G, Veit G, Sun M, Klement JF, et al. Type XIV Collagen Regulates Fibrillogenesis: PREMATURE COLLAGEN FIBRIL GROWTH AND TISSUE DYSFUNCTION IN NULL MICE. *J Biol Chem*. 2009; 284: 8427-8438.
- 9) Liu X, Wu H, Byrne M, Krane S, Jaenisch R. Type III collagen is crucial for collagen I fibrillogenesis and for normal cardiovascular development. *Proc Natl Acad Sci U S A*. 1997; 94: 1852-1856.
- 10) Wenstrup RJ, Florer JB, Brunskill EW, Bell SM, Chervoneva I, Birk DE. Type V Collagen Controls the Initiation of Collagen Fibril Assembly. *J Biol Chem*. 2004; 279: 53331-53337.
- 11) Park AC, Phillips CL, Pfeiffer FM, Roenneburg DA, Kernien JF, Adams SM, et al. Homozygosity and Heterozygosity for Null Col5a2 Alleles Produce Embryonic Lethality and a Novel Classic Ehlers-Danlos Syndrome-Related Phenotype. *Am J Pathol*. 2015; 185: 2000-2011.
- 12) Gelse K. Collagens—structure, function, and biosynthesis. *Adv Drug Deliv Rev*. 2003; 55: 1531-

1546.

- 13) Garofalo S, Vuorio E, Vaughan J, Lozano G, Mayneo R, Ellard J, et al. Reduced amounts of cartilage collagen fibrils and growth plate anomalies in transgenic mice harboring a glycine-to-cysteine mutation in the mouse type II procollagen α -chain gene. *Proc Natl Acad Sci U S A*. 1991; 88: 9648-9652.
- 14) Li SW, Prockop DJ, Helminen H, Fässler R, Lapveteläinen T, Kiraly K, et al. Transgenic mice with targeted inactivation of the Col2 alpha 1 gene for collagen II develop a skeleton with membranous and periosteal bone but no endochondral bone. *Genes Dev*. 1995; 9: 2821-2830.
- 15) Li Y, Lacerda DA, Warman ML, Beier DR, Yoshioka H, Ninomiya Y, et al. A fibrillar collagen gene, Col11a1, is essential for skeletal morphogenesis. *Cell*. 1995; 80: 423-430.
- 16) Li SW, Takanosu M, Arita M, Bao Y, Ren ZX, Maier A, et al. Targeted disruption of Col11a2 produces a mild cartilage phenotype in transgenic mice: Comparison with the human disorder otospondylomegapiphyseal dysplasia (OSMED). *Dev Dyn*. 2001; 222: 141-152.
- 17) Fässler R, Schneegelsberg PN, Dausman J, Shinya T, Muragaki Y, McCarthy MT, et al. Mice lacking alpha 1 (IX) collagen develop noninflammatory degenerative joint disease. *Proc Natl Acad Sci U S A*. 1994; 91: 5070-5074.
- 18) Nakata K, Ono K, Miyazaki J, Olsen BR, Muragaki Y, Adachi E, et al. Osteoarthritis associated with mild chondrodysplasia in transgenic mice expressing alpha 1(IX) collagen chains with a central deletion. *Proc Natl Acad Sci U S A*. 1993; 90: 2870-2874.
- 19) Muragaki Y, Mariman EC, van Beersum SE, Perälä M, van Mourik JB, Warman ML, et al. A mutation in the gene encoding the alpha 2 chain of the fibril-associated collagen IX, COL9A2, causes multiple epiphyseal dysplasia (EDM2). *Nat Genet*. 1996; 12: 103-105.
- 20) Yang L, Tsang KY, Tang HC, Chan D, Cheah KSE. Hypertrophic chondrocytes can become osteoblasts and osteocytes in endochondral bone formation. *Proc Natl Acad Sci U S A*. 2014; 111: 12097-12102.
- 21) Gress CJ, Jacenko O. Growth plate compressions and altered hematopoiesis in collagen X null mice. *J Cell Biol*. 2000; 149: 983-993.
- 22) Boot-Handford RP, Tuckwell DS, Plumb DA, Rock CF, Poulson R. A novel and highly conserved collagen (pro(alpha)1(XXVII)) with a unique expression pattern and unusual molecular characteristics establishes a new clade within the vertebrate fibrillar collagen family. *J Biol Chem*. 2003; 278: 31067-31077.
- 23) Koch M, Laub F, Zhou P, Hahn RA, Tanaka S, Burgeson RE, et al. Collagen XXIV, a vertebrate fibrillar collagen with structural features of invertebrate collagens: selective expression in developing cornea and bone. *J Biol Chem*. 2003; 278: 43236-43244.
- 24) Plumb DA, Ferrara L, Torbica T, Knowles L, Jr AM, Kadler KE, et al. Collagen XXVII Organises the Pericellular Matrix in the Growth Plate. *PLOS ONE*. 2011; 6: e29422.

- 25) Sundaramoorthy M, Meiyappan M, Todd P, Hudson BG. Crystal Structure of NC1 Domains: STRUCTURAL BASIS FOR TYPE IV COLLAGEN ASSEMBLY IN BASEMENT MEMBRANES. *J Biol Chem.* 2002; 277: 31142-31153.
- 26) Gould DB, Phalan FC, Breedveld GJ, van Mil SE, Smith RS, Schimenti JC, et al. Mutations in Col4a1 cause perinatal cerebral hemorrhage and porencephaly. *Science.* 2005; 308: 1167-1171.
- 27) Labelle-Dumais C, Dilworth DJ, Harrington EP, de Leau M, Lyons D, Kabaeva, Z et al. COL4A1 mutations cause ocular dysgenesis, neuronal localization defects, and myopathy in mice and Walker-Warburg syndrome in humans. *PLoS Genet.* 2011; 7: e1002062.
- 28) Favor J, Gloeckner CJ, Janik D, Klempt M, Neuhäuser-Klaus A, Pretsch W, et al. Type IV procollagen missense mutations associated with defects of the eye, vascular stability, the brain, kidney function and embryonic or postnatal viability in the mouse, *Mus musculus*: an extension of the Col4a1 allelic series and the identification of the first two Col4a2 mutant alleles. *Genetics.* 2007; 175: 725-736.
- 29) Cosgrove D, Meehan DT, Grunkemeyer JA, Kornak JM, Sayers R, Hunter WJ, et al. Collagen COL4A3 knockout: a mouse model for autosomal Alport syndrome. *Genes Dev.* 1996; 10: 2981-2992.
- 30) Komori T, Ono M, Hara ES, Ueda J, Nguyen HTT, Nguyen HT, et al. Type IV collagen $\alpha 6$ chain is a regulator of keratin 10 in keratinization of oral mucosal epithelium. *Sci Rep.* 2018; 8: 2612.
- 31) Nishie W, Sawamura D, Goto M, Ito K, Shibaki A, McMillan JR, et al. Humanization of autoantigen. *Nat Med.* 2007; 13: 378-383.
- 32) Tanimura S, Tadokoro Y, Inomata K, Binh NT, Nishie W, Yamazaki S, et al. Hair Follicle Stem Cells Provide a Functional Niche for Melanocyte Stem Cells. *Cell Stem Cell.* 2011; 8: 177-187.
- 33) Matsumura H, Mohri Y, Binh NT, Morinaga H, Fukuda M, Ito M, et al. Hair follicle aging is driven by transepidermal elimination of stem cells via COL17A1 proteolysis. *Science.* 2016; 351: 575-589.
- 34) Heinonen S, Männikkö M, Klement JF, Whitaker-Menezes D, Murphy GF, Uitto J. Targeted inactivation of the type VII collagen gene (Col7a1) in mice results in severe blistering phenotype: a model for recessive dystrophic epidermolysis bullosa. *J Cell Sci.* 1999; 112: 3641-3648.
- 35) Fritsch A, Loeckermann S, Kern JS, Braun A, Bösl MR, Bley TA, et al. A hypomorphic mouse model of dystrophic epidermolysis bullosa reveals mechanisms of disease and response to fibroblast therapy. *J Clin Invest.* 2008; 118: 1669-1679.
- 36) Woodley DT, Keene DR, Atha T, Huang Y, Lipman K, Li W, et al. Injection of recombinant human type VII collagen restores collagen function in dystrophic epidermolysis bullosa. *Nat Med.* 2004; 10: 693-695.
- 37) Kvist A-P, Latvanlehto A, Sund M, Eklund L, Väisänen T, Hägg P et al. Lack of Cytosolic and Transmembrane Domains of Type XIII Collagen Results in Progressive Myopathy. *Am J Pathol.* 2001; 159: 1581-1592.

- 38) Härönen H, Zainul Z, Tu H, Naumenko N, Sormunen R, Miinalainen I, et al. Collagen XIII secures pre- and postsynaptic integrity of the neuromuscular synapse. *Hum Mol Genet.* 2017; 26: 2076-2090.
- 39) Tanaka T, Wakabayashi T, Oizumi H, Nishio S, Sato T, Harada A, et al. CLAC-P/Collagen Type XXV Is Required for the Intramuscular Innervation of Motoneurons during Neuromuscular Development. *J Neurosci.* 2014; 34: 1370-1379.
- 40) Sumiyoshi H, Mor N, Lee SY, Doty S, Henderson S, Tanaka S, et al. Esophageal muscle physiology and morphogenesis require assembly of a collagen XIX-rich basement membrane zone. *J Cell Biol.* 2004; 166: 591-600.
- 41) Su J, Chen J, Lippold K, Monavarfeshani A, Carrillo GL, Jenkins R, et al. Collagen-derived matricryptins promote inhibitory nerve terminal formation in the developing neocortex. *J Cell Biol.* 2016; 212: 721-736.
- 42) Charvet B, Guiraud A, Malbouyres M, Zwolanek D, Guillon E, Bretaud S, et al. Knockdown of *col22a1* gene in zebrafish induces a muscular dystrophy by disruption of the myotendinous junction. *Development.* 2013; 140: 4602-4613.
- 43) Irwin WA, Bergamin N, Sabatelli P, Reggiani C, Megighian A, Merlini L, et al. Mitochondrial dysfunction and apoptosis in myopathic mice with collagen VI deficiency. *Nat Genet.* 2003; 35: 367-371.
- 44) Pan TC, Zhang RZ, Markova D, Arita M, Zhang Y, Bogdanovich S, et al. COL6A3 protein deficiency in mice leads to muscle and tendon defects similar to human collagen VI congenital muscular dystrophy. *J Biol Chem.* 2013; 288: 14320-14331.
- 45) Urciuolo A, Quarta M, Morbidoni V, Gattazzo F, Molon S, Grumati P, et al. Collagen VI regulates satellite cell self-renewal and muscle regeneration. *Nat Commun.* 2013; 4: 1964.
- 46) Greenhill NS, Rüger BM, Hasan Q, Davis PF. The alpha1(VIII) and alpha2(VIII) collagen chains form two distinct homotrimeric proteins in vivo. *Matrix Biol.* 2000; 19: 19-28.
- 47) Hopfer U, Fukai N, Hopfer H, Wolf G, Joyce N, Li E, et al. Targeted disruption of *Col8a1* and *Col8a2* genes in mice leads to anterior segment abnormalities in the eye. *FASEB J.* 2005; 19: 1232-1244.
- 48) Fukai N, Eklund L, Marneros AG, Oh SP, Keene DR, Tamarkin L, et al. Lack of collagen XVIII/endostatin results in eye abnormalities. *EMBO J.* 2002; 21: 1535-1544.
- 49) O'Reilly MS, Boehm T, Shing Y, Fukai N, Vasios G, Lane WS, et al. Endostatin: An Endogenous Inhibitor of Angiogenesis and Tumor Growth. *Cell.* 1997; 88: 277-285.

■Chapter 5. Biosynthesis (1): Overview

ISHIKAWA Yoshihiro

Department of Ophthalmology, University of California, San Francisco School of Medicine

- 1) Ishikawa Y, Bachinger HP. A molecular ensemble in the rER for procollagen maturation. *Biochim Biophys Acta* 2013;1833:2479–91.
- 2) Gorres KL, Raines RT. Prolyl 4-hydroxylase. *Crit Rev Biochem Mol Biol* 2010;45:106–24.
- 3) Yamauchi M, Terajima M, Shiiba M. Lysine Hydroxylation and Cross-Linking of Collagen. *Methods Mol Biol* 2019;1934:309–24.
- 4) Montgomery NT, Zientek KD, Pokidysheva EN, Bachinger HP. Post-translational modification of type IV collagen with 3-hydroxyproline affects its interactions with glycoprotein VI and nidogens 1 and 2. *J Biol Chem* 2018;293:5987–99.
- 5) Taga Y, Kusubata M, Ogawa-Goto K, Hattori S. Developmental Stage-dependent Regulation of Prolyl 3-Hydroxylation in Tendon Type I Collagen. *J Biol Chem* 2016;291:837–47.
- 6) Van Dijk FS, Nesbitt IM, Nikkels PG, et al. CRTAP mutations in lethal and severe osteogenesis imperfecta: the importance of combining biochemical and molecular genetic analysis. *Eur J Hum Genet* 2009;17:1560–9.
- 7) Marini JC, Forlino A, Bachinger HP, et al. Osteogenesis imperfecta. *Nat Rev Dis Primers* 2017;3:17052.
- 8) Myllyharju J, Kivirikko KI. Collagens, modifying enzymes and their mutations in humans, flies and worms. *Trends Genet* 2004;20:33–43.
- 9) Koivunen P, Tiainen P, Hyvarinen J, et al. An endoplasmic reticulum transmembrane prolyl 4-hydroxylase is induced by hypoxia and acts on hypoxia-inducible factor alpha. *J Biol Chem* 2007;282:30544–52.
- 10) Holster T, Pakkanen O, Soininen R, et al. Loss of assembly of the main basement membrane collagen, type IV, but not fibril-forming collagens and embryonic death in collagen prolyl 4-hydroxylase I null mice. *J Biol Chem* 2007;282:2512–9.
- 11) Vranka JA, Sakai LY, Bachinger HP. Prolyl 3-hydroxylase 1, enzyme characterization and identification of a novel family of enzymes. *J Biol Chem* 2004;279:23615–21.
- 12) Morello R, Bertin TK, Chen Y, et al. CRTAP is required for prolyl 3-hydroxylation and mutations cause recessive osteogenesis imperfecta. *Cell* 2006;127:291–304.
- 13) Wu J, Zhang W, Xia L, et al. Characterization of PPIB interaction in the P3H1 ternary complex and implications for its pathological mutations. *Cell Mol Life Sci* 2019;76:3899–914.
- 14) Chang W, Barnes AM, Cabral WA, Bodurtha JN, Marini JC. Prolyl 3-hydroxylase 1 and CRTAP are mutually stabilizing in the endoplasmic reticulum collagen prolyl 3-hydroxylation complex. *Hum Mol Genet* 2010;19:223–34.

- 15) Mattoteia D, Chiapparino A, Fumagalli M, De Marco M, De Giorgi F, Negro L, Pinnola A, Faravelli S, Roscioli T, Scietti L, Forneris F. Identification of Regulatory Molecular "Hot Spots" for LH/PLOD Collagen Glycosyltransferase Activity. *Int J Mol Sci.* 2023 Jul 7;24(13):11213.
- 16) Ishikawa Y, Taga Y, Coste T, Tufa SF, Keene DR, Mizuno K, Tournier-Lasserre E, Gould DB. Lysyl hydroxylase 3-mediated post-translational modifications are required for proper biosynthesis of collagen $\alpha 1\alpha 1\alpha 2(IV)$. *J Biol Chem.* 2022 Dec;298(12):102713.
- 17) Piersma B, Bank RA. Collagen cross-linking mediated by lysyl hydroxylase 2: an enzymatic battlefield to combat fibrosis. *Essays Biochem* 2019;63:377-87.
- 18) Gjaltema RA, van der Stoep MM, Boersema M, Bank RA. Disentangling mechanisms involved in collagen pyridinoline cross-linking: The immunophilin FKBP65 is critical for dimerization of lysyl hydroxylase 2. *Proc Natl Acad Sci U S A* 2016;113:7142-7.
- 19) Hyry M, Lantto J, Myllyharju J. Missense mutations that cause Bruck syndrome affect enzymatic activity, folding, and oligomerization of lysyl hydroxylase 2. *J Biol Chem* 2009;284:30917-24.
- 20) Malfait F. Vascular aspects of the Ehlers-Danlos Syndromes. *Matrix Biol* 2018;71-72:380-95.
- 21) Ishikawa Y, Taga Y, Zientek K, Mizuno N, Salo AM, Semenova O, Tufa SF, Keene DR, Holden P, Mizuno K, Gould DB, Myllyharju J, Bächinger HP. Type I and type V procollagen triple helix uses different subsets of the molecular ensemble for lysine posttranslational modifications in the rER. *J Biol Chem.* 2021 Jan-Jun;296:100453.
- 22) Wilhelm D, Wurtz A, Abouelfarah H, Sanchez G, Bui C, Vincourt JB. Tissue-specific collagen hydroxylation at GEP/GDP triplets mediated by P4HA2. *Matrix Biol.* 2023 May;119:141-153
- 23) Salo AM, Rappu P, Koski MK, Karjalainen E, Izzi V, Drushinin K, Miinalainen I, Käpylä J, Heino J, Myllyharju J. Collagen prolyl 4-hydroxylase isoenzymes I and II have sequence specificity towards different X-Pro-Gly triplets. *Matrix Biol.* 2024 Jan;125:73-87.
- 24) Hudson DM, Joeng KS, Werther R, et al. Post-translationally abnormal collagens of prolyl 3-hydroxylase-2 null mice offer a pathobiological mechanism for the high myopia linked to human LEPREL1 mutations. *J Biol Chem* 2015;290:8613-22.
- 25) Hudson DM, Weis M, Eyre DR. Insights on the evolution of prolyl 3-hydroxylation sites from comparative analysis of chicken and *Xenopus* fibrillar collagens. *PLoS One* 2011;6:e19336.
- 26) Pokidysheva E, Zientek KD, Ishikawa Y, et al. Posttranslational modifications in type I collagen from different tissues extracted from wild type and prolyl 3-hydroxylase 1 null mice. *J Biol Chem* 2013;288:24742-52.
- 27) Heard ME, Besio R, Weis M, et al. Sc65-Null Mice Provide Evidence for a Novel Endoplasmic Reticulum Complex Regulating Collagen Lysyl Hydroxylation. *PLoS Genet* 2016;12:e1006002.
- 28) Hudson DM, Weis M, Rai J, et al. P3h3-null and Sc65-null Mice Phenocopy the Collagen Lysine Under-hydroxylation and Cross-linking Abnormality of Ehlers-Danlos Syndrome Type VIA. *J Biol Chem* 2017;292:3877-87.

- 29) Terajima M, Taga Y, Cabral WA, et al. Cyclophilin B control of lysine post-translational modifications of skin type I collagen. *PLoS Genet* 2019;15:e1008196.
- 30) Terajima M, Taga Y, Chen Y, et al. Cyclophilin-B Modulates Collagen Cross-linking by Differentially Affecting Lysine Hydroxylation in the Helical and Telopeptidyl Domains of Tendon Type I Collagen. *J Biol Chem* 2016;291:9501-12.
- 31) Walter P, Ron D. The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 2011;334:1081-6.
- 32) Ishikawa T, Okada T, Ishikawa-Fujiwara T, et al. ATF6alpha/beta-mediated adjustment of ER chaperone levels is essential for development of the notochord in medaka fish. *Mol Biol Cell* 2013;24:1387-95.
- 33) Ishikawa T, Toyama T, Nakamura Y, et al. UPR transducer BFF2H7 allows export of type II collagen in a cargo- and developmental stage-specific manner. *J Cell Biol* 2017;216:1761-74.
- 34) Symoens S, Malfait F, D'Hondt S, et al. Deficiency for the ER-stress transducer OASIS causes severe recessive osteogenesis imperfecta in humans. *Orphanet J Rare Dis* 2013;8:154.
- 35) Boudko SP, Engel J, Bachinger HP. The crucial role of trimerization domains in collagen folding. *Int J Biochem Cell Biol* 2012;44:21-32.
- 36) Persikov AV, Ramshaw JA, Kirkpatrick A, Brodsky B. Amino acid propensities for the collagen triple-helix. *Biochemistry* 2000;39:14960-7.
- 37) Ishikawa Y, Boudko S, Bachinger HP. Ziploc-ing the structure: Triple helix formation is coordinated by rough endoplasmic reticulum resident PPIases. *Biochim Biophys Acta* 2015;1850:1983-93.
- 38) Rulten SL, Kinloch RA, Tateossian H, Robinson C, Gettins L, Kay JE. The human FK506-binding proteins: characterization of human FKBP19. *Mamm Genome* 2006;17:322-31.
- 39) Zeng B, MacDonald JR, Bann JG, et al. Chicken FK506-binding protein, FKBP65, a member of the FKBP family of peptidylprolyl cis-trans isomerases, is only partially inhibited by FK506. *Biochem J* 1998;330 (Pt 1):109-14.
- 40) Ishikawa Y, Vranka J, Wirz J, Nagata K, Bachinger HP. The rough endoplasmic reticulum-resident FK506-binding protein FKBP65 is a molecular chaperone that interacts with collagens. *J Biol Chem* 2008;283:31584-90.
- 41) Gidalevitz T, Stevens F, Argon Y. Orchestration of secretory protein folding by ER chaperones. *Biochim Biophys Acta* 2013;1833:2410-24.
- 42) Ishikawa Y, Vranka JA, Boudko SP, et al. Mutation in cyclophilin B that causes hyperelastosis cutis in American Quarter Horse does not affect peptidylprolyl cis-trans isomerase activity but shows altered cyclophilin B-protein interactions and affects collagen folding. *J Biol Chem* 2012;287:22253-65.
- 43) Cabral WA, Chang W, Barnes AM, et al. Prolyl 3-hydroxylase 1 deficiency causes a recessive

- metabolic bone disorder resembling lethal/severe osteogenesis imperfecta. *Nat Genet* 2007;39:359–65.
- 44) Friedman J, Weissman I. Two cytoplasmic candidates for immunophilin action are revealed by affinity for a new cyclophilin: one in the presence and one in the absence of CsA. *Cell* 1991;66:799–806.
 - 45) Chapman DC, Stocki P, Williams DB. Cyclophilin C Participates in the US2-Mediated Degradation of Major Histocompatibility Complex Class I Molecules. *PLoS One* 2015;10:e0145458.
 - 46) Stocki P, Chapman DC, Beach LA, Williams DB. Depletion of cyclophilins B and C leads to dysregulation of endoplasmic reticulum redox homeostasis. *J Biol Chem* 2014;289:23086–96.
 - 47) Ito S, Nagata K. Biology of Hsp47 (Serpin H1), a collagen-specific molecular chaperone. *Semin Cell Dev Biol* 2017;62:142–51.
 - 48) Nagai N, Hosokawa M, Itohara S, et al. Embryonic lethality of molecular chaperone hsp47 knockout mice is associated with defects in collagen biosynthesis. *J Cell Biol* 2000;150:1499–506.
 - 49) Christiansen HE, Schwarze U, Pyott SM, et al. Homozygosity for a missense mutation in SERPINH1, which encodes the collagen chaperone protein HSP47, results in severe recessive osteogenesis imperfecta. *Am J Hum Genet* 2010;86:389–98.
 - 50) Ono T, Miyazaki T, Ishida Y, Uehata M, Nagata K. Direct in vitro and in vivo evidence for interaction between Hsp47 protein and collagen triple helix. *J Biol Chem* 2012;287:6810–8.
 - 51) Koide T, Asada S, Takahara Y, Nishikawa Y, Nagata K, Kitagawa K. Specific recognition of the collagen triple helix by chaperone HSP47: minimal structural requirement and spatial molecular orientation. *J Biol Chem* 2006;281:3432–8.
 - 52) Koide T, Nishikawa Y, Asada S, et al. Specific recognition of the collagen triple helix by chaperone HSP47. II. The HSP47-binding structural motif in collagens and related proteins. *J Biol Chem* 2006;281:11177–85.
 - 53) Koide T, Takahara Y, Asada S, Nagata K. Xaa-Arg-Gly triplets in the collagen triple helix are dominant binding sites for the molecular chaperone HSP47. *J Biol Chem* 2002;277:6178–82.
 - 54) Ishikawa Y, Holden P, Bachinger HP. Heat shock protein 47 and 65-kDa FK506-binding protein weakly but synergistically interact during collagen folding in the endoplasmic reticulum. *J Biol Chem* 2017;292:17216–24.
 - 55) Ishikawa Y, Mizuno K, Bachinger HP. Ziploc-ing the structure 2.0: Endoplasmic reticulum-resident peptidyl prolyl isomerases show different activities toward hydroxyproline. *J Biol Chem* 2017;292:9273–82.
 - 56) Ishikawa Y, Bachinger HP. A substrate preference for the rough endoplasmic reticulum resident protein FKBP22 during collagen biosynthesis. *J Biol Chem* 2014;289:18189–201.
 - 57) Ishikawa Y, Bonna A, Gould DB, Farndale RW. Local Net Charge State of Collagen Triple Helix

- Is a Determinant of FKBP22 Binding to Collagen III. *Int J Mol Sci.* 2023 Oct 13;24(20):15156.
- 58) Baumann M, Giunta C, Krabichler B, et al. Mutations in FKBP14 cause a variant of Ehlers–Danlos syndrome with progressive kyphoscoliosis, myopathy, and hearing loss. *Am J Hum Genet* 2012;90:201–16.
- 59) Giunta C, Baumann M, Fauth C, et al. A cohort of 17 patients with kyphoscoliotic Ehlers–Danlos syndrome caused by biallelic mutations in FKBP14: expansion of the clinical and mutational spectrum and description of the natural history. *Genet Med* 2018;20:42–54.
- 60) Kalamajski S, Oldberg A. The role of small leucine-rich proteoglycans in collagen fibrillogenesis. *Matrix Biol* 2010;29:248–53.
- 61) Ishikawa Y, Rubin K, Bachinger HP, Kalamajski S. The endoplasmic reticulum-resident collagen chaperone Hsp47 interacts with and promotes the secretion of decorin, fibromodulin, and lumican. *J Biol Chem* 2018;293:13707–16.
- 62) Chen J, Wang S, Zhang Z, Richards CI, Xu R. Heat shock protein 47 (HSP47) binds to discoidin domain-containing receptor 2 (DDR2) and regulates its protein stability. *J Biol Chem* 2019;294:16846–54.
- 63) Ishikawa Y, Ito S, Nagata K, Sakai LY, Bachinger HP. Intracellular mechanisms of molecular recognition and sorting for transport of large extracellular matrix molecules. *Proc Natl Acad Sci U S A* 2016;113:E6036–E44.
- 64) Sepulveda D, Rojas-Rivera D, Rodriguez DA, et al. Interactome Screening Identifies the ER Luminal Chaperone Hsp47 as a Regulator of the Unfolded Protein Response Transducer IRE1alpha. *Mol Cell* 2018;69:238–52 e7.
- 65) Macdonald JR, Bachinger HP. HSP47 binds cooperatively to triple helical type I collagen but has little effect on the thermal stability or rate of refolding. *J Biol Chem* 2001;276:25399–403.
- 66) Oecal S, Socher E, Uthoff M, et al. The pH-dependent Client Release from the Collagen-specific Chaperone HSP47 Is Triggered by a Tandem Histidine Pair. *J Biol Chem* 2016;291:12612–26.
- 67) Lamande SR, Chessler SD, Golub SB, et al. Endoplasmic reticulum-mediated quality control of type I collagen production by cells from osteogenesis imperfecta patients with mutations in the pro alpha 1 (I) chain carboxyl-terminal propeptide which impair subunit assembly. *J Biol Chem* 1995;270:8642–9.
- 68) Duran I, Martin JH, Weis MA, et al. A Chaperone Complex Formed by HSP47, FKBP65, and BiP Modulates Telopeptide Lysyl Hydroxylation of Type I Procollagen. *J Bone Miner Res* 2017;32:1309–19.
- 69) Gomez-Navarro N, Miller E. Protein sorting at the ER–Golgi interface. *J Cell Biol* 2016;215:769–78.
- 70) McCaughey J, Stephens DJ. ER-to-Golgi Transport: A Sizeable Problem. *Trends Cell Biol*

2019;29:940–53.

- 71) Saito K, Chen M, Bard F, et al. TANGO1 facilitates cargo loading at endoplasmic reticulum exit sites. *Cell* 2009;136:891–902.
- 72) Raote I, Ortega Bellido M, Pirozzi M, et al. TANGO1 assembles into rings around COPII coats at ER exit sites. *J Cell Biol* 2017;216:901–9.
- 73) Ma W, Goldberg J. TANGO1/cTAGE5 receptor as a polyvalent template for assembly of large COPII coats. *Proc Natl Acad Sci U S A* 2016;113:10061–6.
- 74) Gorur A, Yuan L, Kenny SJ, Baba S, Xu K, Schekman R. COPII-coated membranes function as transport carriers of intracellular procollagen I. *J Cell Biol* 2017;216:1745–59.
- 75) McCaughey J, Stevenson NL, Cross S, Stephens DJ. ER-to-Golgi trafficking of procollagen in the absence of large carriers. *J Cell Biol* 2019;218:929–48.
- 76) Omari S, Makareeva E, Roberts-Pilgrim A, et al. Noncanonical autophagy at ER exit sites regulates procollagen turnover. *Proc Natl Acad Sci U S A* 2018;115:E10099–E108.
- 77) Forrester A, De Leonibus C, Grumati P, Fasana E, Piemontese M, Staiano L, Fregno I, Raimondi A, Marazza A, Bruno G, Iavazzo M, Intartaglia D, Seczynska M, van Anken E, Conte I, De Matteis MA, Dikic I, Molinari M, Settembre C. A selective ER-phagy exerts procollagen quality control via a Calnexin-FAM134B complex. *EMBO J*. 2019 Jan 15;38(2):e99847
- 78) Raote I, Malhotra V. Protein transport by vesicles and tunnels. *J Cell Biol* 2019;218:737–9.
- 79) Reynolds HM, Zhang L, Tran DT, Ten Hagen KG. Tango1 coordinates the formation of endoplasmic reticulum/Golgi docking sites to mediate secretory granule formation. *J Biol Chem*. 2019;294(51):19498–19510.
- 80) Raote I, Ernst MA2, Campelo F, Rothman EJ, Pincet F, Malhotra V. TANGO1 membrane helices create a lipid diffusion barrier at curved membranes. *Elife* 2020;9:e57822
- 81) Rezaei N, Lyons A, Forde NR. Environmentally Controlled Curvature of Single Collagen Proteins. *Biophys J* 2018;115:1457–69.
- 82) Al-Shaer A, Lyons A, Ishikawa Y, Hudson BG, Boudko SP, Forde NR. Sequence-dependent mechanics of collagen reflect its structural and functional organization. *Biophys J*. 2021 Sep 21;120(18):4013–4028.
- 83) Raote I, Ortega-Bellido M, Santos AJ, et al. TANGO1 builds a machine for collagen export by recruiting and spatially organizing COPII, tethers and membranes. *Elife* 2018;7.
- 84) Santos AJ, Raote I, Scarpa M, Brouwers N, Malhotra V. TANGO1 recruits ERGIC membranes to the endoplasmic reticulum for procollagen export. *Elife* 2015;4.

■Chapter 6. Biosynthesis (2): Post-translational Modifications

TAGA Yuki

Nippi Research Institute of Biomatrix

- 1) Zou Y, Donkervoort S, Salo AM, Foley AR, Barnes AM, Hu Y, et al. P4HA1 mutations cause a unique congenital disorder of connective tissue involving tendon, bone, muscle and the eye. *Hum Mol Genet.* 2017; 26: 2207–17.
- 2) Cabral WA, Chang W, Barnes AM, Weis M, Scott MA, Leikin S, et al. Prolyl 3-hydroxylase 1 deficiency causes a recessive metabolic bone disorder resembling lethal/severe osteogenesis imperfecta. *Nat Genet.* 2007; 39: 359–65.
- 3) Yeowell HN, Walker LC. Mutations in the lysyl hydroxylase 1 gene that result in enzyme deficiency and the clinical phenotype of Ehlers–Danlos syndrome type VI. *Mol Genet Metab.* 2000; 71: 212–24.
- 4) van der Slot AJ, Zuurmond AM, Bardeel AF, Wijmenga C, Pruijs HE, Sillence DO, et al. Identification of PLOD2 as telopeptide lysyl hydroxylase, an important enzyme in fibrosis. *J Biol Chem.* 2003; 278: 40967–72.
- 5) Salo AM, Cox H, Farndon P, Moss C, Grindulis H, Risteli M, et al. A connective tissue disorder caused by mutations of the lysyl hydroxylase 3 gene. *Am J Hum Genet.* 2008; 83: 495–503.
- 6) Miyatake S, Schneeberger S, Koyama N, Yokochi K, Ohmura K, Shiina M, et al. Biallelic COLGALT1 variants are associated with cerebral small vessel disease. *Ann Neurol.* 2018; 84: 843–53.
- 7) Taga Y, Kusubata M, Ogawa–Goto K, Hattori S. Developmental stage–dependent regulation of prolyl 3-hydroxylation in tendon type I collagen. *J Biol Chem.* 2016; 291: 837–47.
- 8) Holster T, Pakkanen O, Soininen R, Sormunen R, Nokelainen M, Kivirikko KI, et al. Loss of assembly of the main basement membrane collagen, type IV, but not fibril-forming collagens and embryonic death in collagen prolyl 4-hydroxylase I null mice. *J Biol Chem.* 2007; 282: 2512–9.
- 9) Shoulders MD, Raines RT. Collagen structure and stability. *Annu Rev Biochem.* 2009; 78: 929–58.
- 10) Burjanadze TV. Hydroxyproline content and location in relation to collagen thermal stability. *Biopolymers.* 1979; 18: 931–8.
- 11) Rappu P, Salo AM, Myllyharju J, Heino J. Role of prolyl hydroxylation in the molecular interactions of collagens. *Essays Biochem.* 2019; 63: 325–35.
- 12) Ogle JD, Arlinghaus RB, Lgan MA. 3-Hydroxyproline, a new amino acid of collagen. *J Biol Chem.* 1962; 237: 3667–73.
- 13) Morello R, Bertin TK, Chen Y, Hicks J, Tonachini L, Monticone M, et al. CRTAP is required for prolyl 3- hydroxylation and mutations cause recessive osteogenesis imperfecta. *Cell.* 2006; 127: 291–304.
- 14) van Dijk FS, Nesbitt IM, Zwikstra EH, Nikkels PG, Piersma SR, Fratantoni SA, et al. PPIB

- mutations cause severe osteogenesis imperfecta. *Am J Hum Genet.* 2009; 85: 521–7.
- 15) Mordechai S, Gradstein L, Pasanen A, Ofir R, El Amour K, Levy J, et al. High myopia caused by a mutation in *LEPREL1*, encoding prolyl 3-hydroxylase 2. *Am J Hum Genet.* 2011; 89: 438–45.
 - 16) Hudson DM, Weis M, Rai J, Joeng KS, Dimori M, Lee BH, et al. P3h3-null and Sc65-null mice phenocopy the collagen lysine under-hydroxylation and cross-linking abnormality of Ehlers-Danlos syndrome type VIA. *J Biol Chem.* 2017; 292: 3877–87.
 - 17) Hudson DM, Eyre DR. Collagen prolyl 3-hydroxylation: a major role for a minor post-translational modification? *Connect Tissue Res.* 2013; 54: 245–51.
 - 18) Weis MA, Hudson DM, Kim L, Scott M, Wu JJ, Eyre DR. Location of 3-hydroxyproline residues in collagen types I, II, III, and V/XI implies a role in fibril supramolecular assembly. *J Biol Chem.* 2010; 285: 2580–90.
 - 19) Eyre DR, Weis M, Hudson DM, Wu JJ, Kim L. A novel 3-hydroxyproline (3Hyp)-rich motif marks the triple-helical C terminus of tendon type I collagen. *J Biol Chem.* 2011; 286: 7732–6.
 - 20) Montgomery NT, Zientek KD, Pokidysheva EN, Bachinger HP. Post-translational modification of type IV collagen with 3-hydroxyproline affects its interactions with glycoprotein VI and nidogens 1 and 2. *J Biol Chem.* 2018; 293: 5987–99.
 - 21) Mizuno K, Peyton DH, Hayashi T, Engel J, Bachinger HP. Effect of the -Gly-3(S)-hydroxyprolyl-4(R)-hydroxyprolyl- tripeptide unit on the stability of collagen model peptides. *FEBS J.* 2008; 275: 5830–40.
 - 22) Ma F, Sun R, Tremmel DM, Sackett SD, Odorico J, Li L. Large-scale differentiation and site specific discrimination of hydroxyproline isomers by electron transfer/higher-energy collision dissociation (ETHcD) mass spectrometry. *Anal Chem.* 2018; 90: 5857–64.
 - 23) Merl-Pham J, Basak T, Knüppel L, Ramanujam D, Athanason M, Behr J, et al. Quantitative proteomic profiling of extracellular matrix and site-specific collagen post-translational modifications in an in vitro model of lung fibrosis. *Matrix Biol Plus.* 2019; 1: 100005.
 - 24) van Huizen NA, Burgers PC, Saintmont F, Brocorens P, Gerbaux P, Stingl C, et al. Identification of 4-hydroxyproline at the Xaa position in collagen by mass spectrometry. *J Proteome Res.* 2019; 18: 2045–51.
 - 25) Taga Y, Kusubata M, Mizuno K. Quantitative analysis of the positional distribution of hydroxyproline in collagenous Gly-Xaa-Yaa sequences by LC-MS with partial acid hydrolysis and precolumn derivatization. *Anal Chem.* 2020; 92: 8427–34.
 - 26) Myllyla R, Wang C, Heikkinen J, Juffer A, Lampela O, Risteli M, et al. Expanding the lysyl hydroxylase toolbox: new insights into the localization and activities of lysyl hydroxylase 3 (LH3). *J Cell Physiol.* 2007; 212: 323–9.
 - 27) Basak T, Vega-Montoto L, Zimmerman LJ, Tabb DL, Hudson BG, Vanacore RM. Comprehensive characterization of glycosylation and hydroxylation of basement membrane collagen IV by high-

- resolution mass spectrometry. *J Proteome Res.* 2016; 15: 245–58.
- 28) Taga Y, Kusubata M, Ogawa-Goto K, Hattori S. Site-specific quantitative analysis of overglycosylation of collagen in osteogenesis imperfecta using hydrazide chemistry and SILAC. *J Proteome Res.* 2013; 12: 2225–32.
- 29) Brinckmann J, Notbohm H, Tronnier M, Acil Y, Fietzek PP, Schmeller W, et al. Overhydroxylation of lysyl residues is the initial step for altered collagen cross-links and fibril architecture in fibrotic skin. *J Invest Dermatol.* 1999; 113: 617–21.
- 30) Lehmann HW, Wolf E, Roser K, Bodo M, Delling G, Muller PK. Composition and posttranslational modification of individual collagen chains from osteosarcomas and osteofibrous dysplasias. *J Cancer Res Clin Oncol.* 1995; 121: 413–8.
- 31) Bailey AJ, Wotton SF, Sims TJ, Thompson PW. Post-translational modifications in the collagen of human osteoporotic femoral head. *Biochem Biophys Res Commun.* 1992; 185: 801–5.
- 32) Terajima M, Perdivara I, Sricholpech M, Deguchi Y, Pleshko N, Tomer KB, et al. Glycosylation and cross-linking in bone type I collagen. *J Biol Chem.* 2014; 289: 22636–47.
- 33) Torre-Blanco A, Adachi E, Hojima Y, Wootton JA, Minor RR, Prockop DJ. Temperature-induced post-translational over-modification of type I procollagen. Effects of over-modification of the protein on the rate of cleavage by procollagen N-proteinase and on self-assembly of collagen into fibrils. *J Biol Chem.* 1992; 267: 2650–5.
- 34) Sricholpech M, Perdivara I, Yokoyama M, Nagaoka H, Terajima M, Tomer KB, et al. Lysyl hydroxylase 3-mediated glucosylation in type I collagen: molecular loci and biological significance. *J Biol Chem.* 2012; 287: 22998–3009.
- 35) Shiraki M, Kuroda T, Tanaka S, Saito M, Fukunaga M, Nakamura T. Nonenzymatic collagen cross-links induced by glycooxidation (pentosidine) predicts vertebral fractures. *J Bone Miner Metab.* 2008; 26: 93–100.
- 36) Yamamoto M, Yamaguchi T, Yamauchi M, Yano S, Sugimoto T. Serum pentosidine levels are positively associated with the presence of vertebral fractures in postmenopausal women with type 2 diabetes. *J Clin Endocrinol Metab.* 2008; 93: 1013–9.
- 37) Iijima K, Murata M, Takahara H, Irie S, Fujimoto D. Identification of N(omega)-carboxymethylarginine as a novel acid-labile advanced glycation end product in collagen. *Biochem J.* 2000; 347 Pt 1: 23–7.
- 38) Odani H, Iijima K, Nakata M, Miyata S, Kusunoki H, Yasuda Y, et al. Identification of N(omega)-carboxymethylarginine, a new advanced glycation endproduct in serum proteins of diabetic patients: possibility of a new marker of aging and diabetes. *Biochem Biophys Res Commun.* 2001; 285: 1232–6.
- 39) Kinoshita S, Mera K, Ichikawa H, Shimasaki S, Nagai M, Taga Y, et al. N(omega) - (carboxymethyl)arginine is one of the dominant advanced glycation end products in glycated

- collagens and mouse tissues. *Oxid Med Cell Longev*. 2019; 2019: 9073451.
- 40) Dunn JA, McCance DR, Thorpe SR, Lyons TJ, Baynes JW. Age-dependent accumulation of N epsilon-(carboxymethyl)lysine and N epsilon-(carboxymethyl)hydroxylysine in human skin collagen. *Biochemistry*. 1991; 30: 1205-10.
- 41) Gorisse L, Pietrement C, Vuiblet V, Schmelzer CE, Kohler M, Duca L, et al. Protein carbamylation is a hallmark of aging. *Proc Natl Acad Sci U S A*. 2016; 113: 1191-6.
- 42) Pietrement C, Gorisse L, Jaisson S, Gillery P. Chronic increase of urea leads to carbamylated proteins accumulation in tissues in a mouse model of CKD. *PLoS One*. 2013; 8: e82506.
- 43) Verbrugge FH, Tang WH, Hazen SL. Protein carbamylation and cardiovascular disease. *Kidney Int*. 2015; 88: 474-8.
- 44) Shi J, Knevel R, Suwannalai P, van der Linden MP, Janssen GM, van Veelen PA, et al. Autoantibodies recognizing carbamylated proteins are present in sera of patients with rheumatoid arthritis and predict joint damage. *Proc Natl Acad Sci U S A*. 2011; 108: 17372-7.
- 45) Koeth RA, Kalantar-Zadeh K, Wang Z, Fu X, Tang WH, Hazen SL. Protein carbamylation predicts mortality in ESRD. *J Am Soc Nephrol*. 2013; 24: 853-61.
- 46) Wang Z, Nicholls SJ, Rodriguez ER, Kumm O, Horkko S, Barnard J, et al. Protein carbamylation links inflammation, smoking, uremia and atherogenesis. *Nat Med*. 2007; 13: 1176-84.
- 47) Jaisson S, Lorimier S, Ricard-Blum S, Sockalingum GD, Delevallee-Forte C, Kegelaer G, et al. Impact of carbamylation on type I collagen conformational structure and its ability to activate human polymorphonuclear neutrophils. *Chem Biol*. 2006; 13: 149-59.
- 48) Taga Y, Tanaka K, Hamada C, Kusubata M, Ogawa-Goto K, Hattori S. Hydroxyhomocitrulline is a collagen-specific carbamylation mark that affects cross-link formation. *Cell Chem Biol*. 2017; 24: 1276-84.
- 49) Hudson DM, Archer M, King KB, Eyre DR. Glycation of type I collagen selectively targets the same helical domain lysine sites as lysyl oxidase-mediated cross-linking. *J Biol Chem*. 2018; 293: 15620-7.

■Chapter 7. Biosynthesis (3): Cross-linking

YAMAUCHI Mitsuo, TERAJIMA Masahiko

University of North Carolina at Chapel Hill

Division, Oral and Craniofacial Health Sciences

- 1) Ricard-Blum S. The collagen family. *Cold Spring Harb Perspect Biol.* 2011 Jan;3(1):a004978. PubMed PMID: 21421911. Pubmed Central PMCID: 3003457.
- 2) Narayanan AS, Siegel RC, Martin GR. On the inhibition of lysyl oxidase by α -aminopropionitrile. *Biochem Biophys Res Commun.* 1972;46(2):745-51.
- 3) Tanzer LM, Housley T, Berube L, Fairweather R, Franzblau C, Gallop, M. Paul. Structure of Two Histidine-containing Cross-Links from Collagen. *J Biol Chem.* 1973;248(2):393-402.
- 4) Robins SP, Bailey AJ. The chemistry of the collagen cross-links. The characterization of fraction C, a possible artifact produced during the reduction of collagen fibres with borohydride. *Biochem J.* 1973;135:657-65.
- 5) Bernstein HP, Mechanic LG. A natural histidine-based imminium cross-link in collagen and its location. *J Biol Chem* 1980;255(21):10414-22.
- 6) Eyre DR, Weis M, Rai J. Analyses of lysine aldehyde cross-linking in collagen reveal that the mature cross-link histidinohydroxylysinoxorleucine is an artifact. *The Journal of biological chemistry.* 2019 Apr 19;294(16):6578-90. PubMed PMID: 30733334. Pubmed Central PMCID: 6484114.
- 7) Yamauchi M, Taga Y, Terajima M. Analyses of lysine aldehyde cross-linking in collagen reveal that the mature cross-link histidinohydroxylysinoxorleucine is an artifact. *The Journal of biological chemistry.* 2019 Sep 20;294(38):14163. PubMed PMID: 31541027. Pubmed Central PMCID: 6755810 this article.
- 8) Housley T, Tanzer LM, Henson E, Gallop MP. Collagen crosslinking: isolation of hydroxyaldol-histidine, a naturally-occurring crosslink. *Biochem Biophys Res Commun* 1975;67(2):824-30.
- 9) Yamauchi M, London RE, Guenat C, Hashimoto F, Mechanic GL. Structure and formation of a stable histidine-based trifunctional cross-link in skin collagen. *J Biol Chem.* 1987;262:11428-34.
- 10) Fujimoto D. Isolation and characterization of a fluorescent material in bovine achilles tendon collagen. *Biochem Biophys Res Commun.* 1977;76(4):1124-9.
- 11) Kuboki Y, Takagi T, Sasaki S, Saito S, G.L. M. Comparative collagen biochemistry of bovine periodontium, gingiva, and dental pulp. *J Dent Res.* 1981;60:159-63.
- 12) Kuboki Y, Tsuzaki M, Sasaki S, Liu CF, Mechanic GL. Location of the intermolecular cross-links in bovine dentin collagen, solubilization with trypsin and isolation of cross-link peptides containing dihydroxylysinoxorleucine and pyridinoline. *Biochem Biophys Res Commun.* 1981;102(1):119-26.
- 13) Yamauchi M, Katz EP, Mechanic GL. Intermolecular cross-linking and stereospecific molecular

- packing in type I collagen fibrils of the periodontal ligament. *Biochemistry*. 1986;25:4907–13.
- 14) Otsubo K, Katz EP, Mechanic GL, Yamauchi M. Cross-linking connectivity in bone collagen fibrils: the COOH-terminal locus of free aldehyde. *Biochemistry*. 1992;31(2):396–402.
 - 15) Yamauchi M, Katz EP. The post-translational chemistry and molecular packing of mineralizing tendon collagens. *Connect Tissue Res*. 1993;29(2):81–98.
 - 16) Fujii K, Tanzer ML. Age-related changes in the reducible crosslinks of human tendon collagen. *FEBS Lett*. 1974;43(3):300–2.
 - 17) Fujii K, Tanzer ML. Osteogenesis imperfecta: biochemical studies of bone collagen. *Clin Orthop Relat Res*. 1977;124:271–7.
 - 18) Saito M, Marumo K. Collagen cross-links as a determinant of bone quality: a possible explanation for bone fragility in aging, osteoporosis, and diabetes mellitus. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2010 Feb;21(2):195–214. PubMed PMID: 19760059.
 - 19) Fujimoto D, Moriguchi T, Ishida T, Hayashi H. The structure of pyridinoline, a collagen crosslink. *Biochem Biophys Res Commun*. 1978;84(1):52–7.
 - 20) Elsdon DF, Light ND, Bailey AJ. An investigation of pyridinoline, and putative collagen cross-link. *Biochem J*. 1980;185(2):531–4.
 - 21) Eyre RD. Cross-linking in collagen and elastin. *Ann Rev Biochem*. 1984;53:717–48.
 - 22) Yamauchi M, Sricholpech M. Lysine post-translational modifications of collagen. *Essays Biochem*. 2012;52:113–33. PubMed PMID: 22708567. Epub 2012/06/20. eng.
 - 23) Eyre DR, Weis MA. Bone Collagen: New Clues to Its Mineralization Mechanism from Recessive Osteogenesis Imperfecta. *Calcified tissue international*. 2013 Mar 19;93:338–47. PubMed PMID: 23508630. Epub 2013/03/20. Eng.
 - 24) Gjaltema RA, Bank RA. Molecular insights into prolyl and lysyl hydroxylation of fibrillar collagens in health and disease. *Crit Rev Biochem Mol Biol*. 2017 Feb;52(1):74–95. PubMed PMID: 28006962.
 - 25) Saito M, Marumo K. Effects of Collagen Crosslinking on Bone Material Properties in Health and Disease. *Calcified tissue international*. 2015 Sep;97(3):242–61. PubMed PMID: 25791570.
 - 26) Ruiz HH, Ramasamy R, Schmidt AM. Advanced Glycation End Products: Building on the Concept of the “Common Soil” in Metabolic Disease. *Endocrinology*. 2020 Jan 1;161(1). PubMed PMID: 31638645.
 - 27) Mechanic GL, Katz EP, Henmi M, Noyes C, Yamauchi M. Locus of a histidine-based, stable trifunctional, helix to helix collagen cross-link: Stereospecific collagen structure of Type I skin fibrils. *Biochemistry*. 1987;26:3500–9.
 - 28) Yamauchi M, Woodley DT, G.L. M. Aging and Cross-linking of Skin Collagen. *Biochem Biophys Res Commun*. 1988;152:893–903.

- 29) Yamauchi M, Prisyanyh P, Haque Z, Woodley DT. Collagen cross-linking in sun-exposed and unexposed sites of aged human skin. *J Invest Dermatol*. 1991;97(5):938-41.
- 30) Kalamajski S, Liu C, Tillgren V, Rubin K, Oldberg A, Rai J, et al. Increased C-telopeptide cross-linking of tendon type I collagen in fibromodulin-deficient mice. *The Journal of biological chemistry*. 2014 Jul 4;289(27):18873-9. PubMed PMID: 24849606. Pubmed Central PMCID: 4081928.
- 31) Terajima M, Taga Y, Cabral WA, Liu Y, Nagasawa M, Sumida N, et al. Cyclophilin B control of lysine post-translational modifications of skin type I collagen. *PLoS Genet*. 2019 Jun 7;15(6):e1008196. PubMed PMID: 31173582.
- 32) Eyre DR, Oguchi H. The hydroxypyridinium crosslinks of skeletal collagens: their measurement, properties and a proposed pathway of formation. *Biochem Biophys Res Commun*. 1980;92(2):403-10.
- 33) Sakura S, Fujimoto D, Sakamoto K, Mizuno A, Motegi K. Photolysis of pyridinoline, a cross-linking amino acid of collagen, by ultraviolet light. *Can J Biochem*. 1982;60(5):525-9.
- 34) Kuypers R, Tyler M, Kurth LB, Jenkins ID, Horgan DJ. Identification of the loci of the collagen-associated Ehrlich chromogen in type I collagen confirms its role as a trivalent cross-link. *Biochem J*. 1992;283:129-36.
- 35) Fujimoto D, Suzuki M, Uchiyama A, Miyamoto S, Inoue T. Analysis of pyridinoline, a cross-linking compound of collagen fibers, in human urine. *J Biochem*. 1983;94(4):1133 - 1136.
- 36) Eyre DR, Weis MA, Wu JJ. Maturation of collagen Ketoimine cross-links by an alternative mechanism to pyridinoline formation in cartilage. *The Journal of biological chemistry*. 2010 May 28;285(22):16675-82. PubMed PMID: 20363745. Pubmed Central PMCID: 2878069. Epub 2010/04/07. eng.
- 37) Uzawa K, Grzesik JW, Nishiura T, Kuzetsov AS, Robey GP, Brenner AD, et al. Differential Expression of Human Lysyl Hydroxylase Genes, Lysine Hydroxylation, and Cross-Linking of Type I Collagen During Osteoblastic Differentiation In Vitro. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 1999;14(8):1272-80.
- 38) Bank AR, Robins PS, Wijmenga C, Breslau-Siderius JL, Bardoel FJA, Van der Sluijs AH, et al. Defective collagen crosslinking in bone, but not in ligament or cartilage, in Bruck syndrome: Indications for a bone-specific telopeptide lysyl hydroxylase on chromosome 17. *Proc Natl Acad Sci USA*. 1999;96(3):1054-8.
- 39) Mercer DK, Nicol PF, Kimbembe C, Robins SP. Identification, expression, and tissue distribution of the three rat lysyl hydroxylase isoforms. *Biochemical and Biophysical Research Communications*. 2003;307(4):803-9.
- 40) Pornprasertsuk S, Duarte WR, Mochida Y, Yamauchi M. Lysyl hydroxylase-2b directs collagen cross-linking pathways in MC3T3-E1 cells. *Journal of bone and mineral research : the official*

journal of the American Society for Bone and Mineral Research. 2004 Aug;19(8):1349–55. PubMed PMID: 15231023.

- 41) Pornprasertsuk S, Duarte WR, Mochida Y, Yamauchi M. Overexpression of lysyl hydroxylase-2b leads to defective collagen fibrillogenesis and matrix mineralization. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2005 Jan;20(1):81–7. PubMed PMID: 15619673. Epub 2004/12/28. eng.
- 42) Sricholpech M, Perdivara I, Nagaoka H, Yokoyama M, Tomer KB, Yamauchi M. Lysyl hydroxylase 3 glucosylates galactosylhydroxylysine residues in type I collagen in osteoblast culture. *The Journal of biological chemistry*. 2011 Mar 18;286(11):8846–56. PubMed PMID: 21220425. Pubmed Central PMCID: 3058983. Epub 2011/01/12. eng.
- 43) Sricholpech M, Perdivara I, Yokoyama M, Nagaoka H, Terajima M, Tomer KB, et al. Lysyl Hydroxylase 3-mediated Glucosylation in Type I Collagen: MOLECULAR LOCI AND BIOLOGICAL SIGNIFICANCE. *Journal of Biological Chemistry*. 2012;287(27):22998–3009.
- 44) Terajima M, Taga Y, Sricholpech M, Kayashima Y, Sumida N, Maeda N, et al. Role of Glycosyltransferase 25 Domain 1 in Type I Collagen Glycosylation and Molecular Phenotypes. *Biochemistry*. 2019 Dec 17;58(50):5040–51. PubMed PMID: 31726007.
- 45) Terajima M, Perdivara I, Sricholpech M, Deguchi Y, Pleshko N, Tomer KB, et al. Glycosylation and cross-linking in bone type I collagen. *The Journal of biological chemistry*. 2014 Aug 15;289(33):22636–47. PubMed PMID: 24958722. Pubmed Central PMCID: 4132771.
- 46) Heard ME, Besio R, Weis M, Rai J, Hudson DM, Dimori M, et al. Sc65-Null Mice Provide Evidence for a Novel Endoplasmic Reticulum Complex Regulating Collagen Lysyl Hydroxylation. *PLoS Genet*. 2016 Apr;12(4):e1006002. PubMed PMID: 27119146. Pubmed Central PMCID: 4847768.
- 47) Duran I, Martin JH, Weis MA, Krejci P, Konik P, Li B, et al. A Chaperone Complex Formed by HSP47, FKBP65, and BiP Modulates Telopeptide Lysyl Hydroxylation of Type I Procollagen. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2017 Feb 08;32(6):1309–19. PubMed PMID: 28177155.
- 48) Gjaltema RA, van der Stoep MM, Boersema M, Bank RA. Disentangling mechanisms involved in collagen pyridinoline cross-linking: The immunophilin FKBP65 is critical for dimerization of lysyl hydroxylase 2. *Proc Natl Acad Sci U S A*. 2016 Jun 13;113(26):7142–7. PubMed PMID: 27298363.
- 49) Schwarze U, Cundy T, Pyott SM, Christiansen HE, Hegde MR, Bank RA, et al. Mutations in FKBP10, which result in Bruck syndrome and recessive forms of osteogenesis imperfecta, inhibit the hydroxylation of telopeptide lysines in bone collagen. *Hum Mol Genet*. 2013 Jan 1;22(1):1–17. PubMed PMID: 22949511. Pubmed Central PMCID: 3606010.
- 50) Cabral WA, Perdivara I, Weis M, Terajima M, Blissett AR, Chang W, et al. Abnormal type I collagen post-translational modification and crosslinking in a cyclophilin B KO mouse model of recessive osteogenesis imperfecta. *PLoS Genet*. 2014 Jun;10(6):e1004465. PubMed PMID: 24968150.

Pubmed Central PMCID: 4072593.

- 51) Terajima M, Taga Y, Chen Y, Cabral WA, Hou-Fu G, Srisawasdi S, et al. Cyclophilin-B modulates collagen cross-linking by differentially affecting lysine hydroxylation in the helical and telopeptidyl domains of tendon type I collagen. *The Journal of biological chemistry*. 2016 Mar 2;291(18):9501–12. PubMed PMID: 26934917.
- 52) Terajima M, Taga Y, Cabral WA, Nagasawa M, Sumida N, Hattori S, et al. Cyclophilin B Deficiency Causes Abnormal Dentin Collagen Matrix. *J Proteome Res*. 2017 Aug 04;16(8):2914–23. PubMed PMID: 28696707.
- 53) Yamauchi M, Noyes C, Kuboki Y, Mechanic GL. Collagen structural microheterogeneity and a possible role for glycosylated hydroxylysine in type I collagen. *Proc Natl Acad Sci U S A*. 1982;79(24):7684–8.
- 54) Gilkes DM, Bajpai S, Wong CC, Chaturvedi P, Hubbi ME, Wirtz D, et al. Procollagen lysyl hydroxylase 2 is essential for hypoxia-induced breast cancer metastasis. *Mol Cancer Res*. 2013 May;11(5):456–66. PubMed PMID: 23378577. Pubmed Central PMCID: 3656974.
- 55) Eisinger-Mathason TS, Zhang M, Qiu Q, Skuli N, Nakazawa MS, Karakasheva T, et al. Hypoxia-dependent modification of collagen networks promotes sarcoma metastasis. *Cancer Discov*. 2013 Oct;3(10):1190–205. PubMed PMID: 23906982. Pubmed Central PMCID: 3822914.
- 56) Chen Y, Terajima M, Yang Y, Sun L, Ahn Y-H, Pankova D, et al. Lysyl hydroxylase 2 induces a collagen cross-link switch in tumor stroma. *The Journal of clinical investigation*. 2015;125(3):1147–62.
- 57) Kirschmann DA, Seftor EA, Nieva DR, Mariano EA, Hendrix MJ. Differentially expressed genes associated with the metastatic phenotype in breast cancer. *Breast Cancer Res Treat*. 1999;55(2):127–36.
- 58) Barker HE, Cox TR, Erler JT. The rationale for targeting the LOX family in cancer. *Nat Rev Cancer*. 2012 Jul 19;12(8):540–52. PubMed PMID: 22810810.
- 59) Cox TR, Bird D, Baker AM, Barker HE, Ho MW, Lang G, et al. LOX-mediated collagen crosslinking is responsible for fibrosis-enhanced metastasis. *Cancer Res*. 2013 Mar 15;73(6):1721–32. PubMed PMID: 23345161. Pubmed Central PMCID: 3672851.
- 60) Levental KR, Yu H, Kass L, Lakins JN, Egeblad M, Erler JT, et al. Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell*. 2009 Nov 25;139(5):891–906. PubMed PMID: 19931152. Pubmed Central PMCID: 2788004.
- 61) Saito T, Uzawa K, Terajima M, Shiiba M, Amelio AL, Tanzawa H, et al. Aberrant Collagen Cross-linking in Human Oral Squamous Cell Carcinoma. *Journal of dental research*. 2019;98(5):517–25.
- 62) Yamauchi M, Barker TH, Gibbons DL, Kurie JM. The fibrotic tumor stroma. *The Journal of clinical investigation*. 2018 Jan 2;128(1):16–25. PubMed PMID: 29293090. Pubmed Central PMCID: 5749516.
- 63) Uzawa K, Marshall, M. K., Katz EP, Tanzawa H, Yeowell HN, Yamauchi M. Altered posttranslational

- modifications of collagen in keloid. *Biochem Biophys Res Commun* 1998;249:652-5.
- 64) Brinckmann J, Notbohm H, Tronnier M, Acil Y, Fietzek PP, Schmeller W, et al. Overhydroxylation of lysyl residues is the initial step for altered collagen cross-links and fibril architecture in fibrotic skin. *J Invest Dermatol.* 1999 Oct;113(4):617-21. PubMed PMID: 10504450.
- 65) van der Slot AJ, Zuurmond AM, Bardoel AF, Wijmenga C, Pruijs HE, Sillence DO, et al. Identification of PLOD2 as telopeptide lysyl hydroxylase, an important enzyme in fibrosis. *The Journal of biological chemistry.* 2003 Oct 17;278(42):40967-72. PubMed PMID: 12881513.
- 66) van der Slot AJ, Zuurmond AM, van den Bogaerdt AJ, Ulrich MMW, Middelkoop E, Boers W, et al. Increased formation of pyridinoline cross-links due to higher telopeptide lysyl hydroxylase levels is a general fibrotic phenomenon. *Matrix Biol* 2004;23:251-7.
- 67) Piersma B, Bank RA. Collagen cross-linking mediated by lysyl hydroxylase 2: an enzymatic battlefield to combat fibrosis. *Essays Biochem.* 2019;63(3):377-38.
- 68) Rosell-Garcia T, Palomo-Alvarez O, Rodriguez-Pascual F. A hierarchical network of hypoxia-inducible factor and SMAD proteins governs procollagen lysyl hydroxylase 2 induction by hypoxia and transforming growth factor beta1. *The Journal of biological chemistry.* 2019 Aug 7. PubMed PMID: 31391253.
- 69) Gilkes DM, Semenza GL, Wirtz D. Hypoxia and the extracellular matrix: drivers of tumour metastasis. *Nat Rev Cancer.* 2014 Jun;14(6):430-9. PubMed PMID: 24827502. Pubmed Central PMCID: 4283800.
- 70) Song Y, Zheng S, Wang J, Long H, Fang L, Wang G, et al. Hypoxia-induced PLOD2 promotes proliferation, migration and invasion via PI3K/Akt signaling in glioma. *Oncotarget.* 2017;8(26):41947-62

■Chapter 8. Collagenase

INADA Masaki

Department of Biotechnology and Life Science, Tokyo University of Agriculture and Technology

- 1) Gross J, Lapiere CM. Collagenolytic activity in amphibian tissues: a tissue culture assay. *Proc Natl Acad Sci U S A*. 1962; 48: 1013–22.
- 2) Eisen AZ, Jeffrey JJ, Gross J. Human skin collagenase. Isolation and mechanism of attack on the collagen molecule. *Biochim Biophys Acta*. 1968; 151: 637–45.
- 3) Nagase H, Woessner Jr JF. Matrix metalloproteinases. *J Biol Chem*. 1999; 274: 21491–4.
- 4) Vu TH, Werb Z. Matrix metalloproteinases: effectors of development and normal physiology. *Genes Dev*. 2000; 14: 2123–33.
- 5) Sternlicht MD, Werb Z. How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol*. 2001; 17: 463–516.
- 6) Puentes XS, López-Otín C. A genomic analysis of rat proteases and protease inhibitors. *Genome Res*. 2004; 14: 609–22.
- 7) Nagase H, Visse R, Murphy G. Structure and function of matrix metalloproteinases and TIMPs. *Cardiovasc Res*. 2006; 69: 562–73.
- 8) Madzharova E, Kastl P, Sabino F, Auf dem Keller U. *Int J Mol Sci*. 2019; 20: 3077.
- 9) Ii M, Yamamoto H, Imai K. *J Electrophoresis*. 2003; 47: 111.
- 10) Brown RA, Hukins DW, Weiss JB, Twose TM. Do mammalian collagenase and DNA restriction endonucleases share a single mechanism for cleavage site recognition? *Biochem Biophys Res Commun*. 1977; 74: 1102–8.
- 11) Mitchell PG, Magna HA, Reeves LM, Lopresti-Morrow LL, Yocum SA, Rosner PJ, et al. Cloning, expression, type II collagenolytic activity of matrix metalloproteinase-13 from human osteoarthritic cartilage. *J Clin Invest*. 1996; 97: 761–8.
- 12) Reboul P, Pelletier JP, Tardif G, Cloutier JM, Martel-Pelletier J. The new collagenase, collagenase-3, is expressed and synthesized by human chondrocytes but not by synoviocytes. A role in osteoarthritis. *J Clin Invest*. 1996; 97: 2011–9.
- 13) Billingham RC, Dahlberg L, Ionescu M, Reiner A, Bourne R, Rorabeck C, et al. Enhanced cleavage of type II collagen by collagenases in osteoarthritic articular cartilage. *J Clin Invest*. 1997; 99: 1534–45.
- 14) Vu TH, Shipley JM, Bergers G, Berger JE, Helms JA, Hanaban D, et al. MMP-9/gelatinase B is a key regulator of growth plate angiogenesis and apoptosis of hypertrophic chondrocytes. *Cell*. 1998; 93: 411–22.
- 15) Zhao W, Byrne MH, Wang Y, Krane SM. Osteocyte and osteoblast apoptosis and excessive bone deposition accompany failure of collagenase cleavage of type I collagen. *J Clin Invest*. 2000;

106: 941–9.

- 16) Zhao W, Byrne MH, Boyce BF, Krane SM. Bone resorption induced by parathyroid hormone is strikingly diminished in collagenase-resistant mutant mice. *J Clin Invest.* 1999; 103: 517–24.
- 17) Sato H, Takino T, Okada Y, Cao J, Shinagawa A, Yamamoto E, et al. Matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nature.* 1994; 370: 61–5.
- 18) Holmbeck K, Bianco P, Caterina J, Yamada S, Kromer M, Kuznetsov SA, et al. MT1MMP-deficient mice develop dwarfism, osteopenia, arthritis, and connective tissue disease due to inadequate collagen turnover. *Cell.* 1999; 99: 81–92.
- 19) Sabeh F, Ota I, Holmbeck K, Birkedal-Hansen H, Soloway P, Balbin M, et al. Tumor cell traffic through the extracellular matrix is controlled by the membrane-anchored collagenase MT1-MMP. *J Cell Biol.* 2004; 167: 769–81.
- 20) Oh J, Takahashi R, Kondo S, Mizoguchi A, Adachi E, Sasahara RM. The membrane-anchored MMP inhibitor RECK is a key regulator of extracellular matrix integrity and angiogenesis. *Cell.* 2001; 107: 789–800.
- 21) Inada M, Wang Y, Byrne MH, Rahman MU, Miyaura C, López-Otín C, et al. Critical roles of collagenase-3 (Mmp13) in development of growth plate cartilage and in endochondral ossification. *Proc Natl Acad Sci U S A.* 2004; 101: 17192–7.
- 22) Krane SM. Possible roles of MMP-13 in skeletal development and bone remodeling: Lessons from mice with MMP-13 null mutation. *Gordon Research Conference, Bones and teeth.* 2000.
- 23) Inada M, Krane SM. Mice with null mutation in collagenase-3 (matrix metalloproteinase [MMP]-13) exhibit altered bone remodeling and increased bone mass. *ASBMR 23th meeting.* 2001; 1052.
- 24) Stickens D, Behonick DJ, Ortega N, Heyer B, Hartenstein B, Yu Y, et al. Altered endochondral bone development in matrix metalloproteinase 13-deficient mice. *Development.* 2004; 131: 5883–95.
- 25) Kosaki N, Takaishi H, Kamekura S, Kimura T, Okada Y, Mingi L, et al. Impaired bone fracture healing in matrix metalloproteinase-13 deficient mice. *Biochem Biophys Res Commun.* 2007; 354: 846–51.
- 26) Deguchi JO, Aikawa E, Liby P, Vachon JR, Inada M, Krane SM, et al. Matrix metalloproteinase-13/collagenase-3 deletion promotes collagen accumulation and organization in mouse atherosclerotic plaques. *Circulation.* 2005; 112: 2708–15.
- 27) Toriseva M, Laato M, Carpén O, Ruohonen ST, Savontaus E, Inada M, et al. MMP-13 regulates growth of wound granulation tissue and modulates gene expression signatures involved in inflammation, proteolysis, and cell viability. *PLoS One.* 2012; 7: e42596.
- 28) Balbín M, Fueyo A, Tester AM, Pendás AM, Pitiot AS, Astudillo A, et al. Loss of collagenase-2 confers increased skin tumor susceptibility to male mice. *Nat Genet.* 2003; 35: 252–7.
- 29) Gutiérrez-Fernández A, Inada M, Balbín M, Fueyo A, Pitiot AS, Astudillo A, et al. Increased

- inflammation delays wound healing in mice deficient in collagenase-2 (MMP-8). *FASEB J.* 2007; 21: 2580-91.
- 30) Inoue K, Mikuni-Takagaki Y, Oikawa K, Itoh T, Inada M, Noguchi T, et al. A crucial role for matrix metalloproteinase 2 in osteocytic canalicular formation and bone metabolism. *J Biol Chem.* 2006; 281: 33814-24.
- 31) Martignetti JA, Aqeel AA, Sewairi WA, Boumah CE, Kambouris M, Mayouf SA, et al. Mutation of the matrix metalloproteinase 2 gene (MMP-2) causes a multicentric osteolysis and arthritis syndrome. *Nat Genet.* 2001; 28: 261-5.
- 32) Kennedy AM, Inada M, Krane SM, Christie PT, Harding B, López-Otín C, et al. MMP13 mutation causes spondyloepimetaphyseal dysplasia, Missouri type (SEMD(MO)). *J Clin Invest.* 2005; 115: 2832-42.
- 33) Nakahara H, Howard L, Thompson EW, Sato H, Seiki M, Yeh Y, et al. Transmembrane/cytoplasmic domain-mediated membrane type 1-matrix metalloprotease docking to invadopodia is required for cell invasion. *Proc Natl Acad Sci USA.* 1997; 94: 7959-64.
- 34) Yu WH, Woessner JF Jr, McNeish JD, Stamenkovic I. CD44 anchors the assembly of matrilysin/MMP-7 with heparin-binding epidermal growth factor precursor and ErbB4 and regulates female reproductive organ remodeling. *Genes Dev.* 2002; 16: 307-23
- 35) Yu Q, Stamenkovic I. Localization of matrix metalloproteinase 9 to the cell surface provides a mechanism for CD44-mediated tumor invasion. *Genes Dev.* 1999; 13: 35-48.
- 36) Yu Q, Stamenkovic I. Cell Surface-Localized Matrix metalloproteinase-9 Proteolytically Activates TGF-beta and Promotes Tumor Invasion and Angiogenesis. *Genes Dev.* 2000; 14: 163-76.
- 37) Brooks PC, Strömblad S, Sanders LC, von Schalscha TL, Aimes RT, Stetler-Stevenson WG, et al. Localization of Matrix Metalloproteinase MMP-2 to the Surface of Invasive Cells by Interaction with Integrin $\alpha v \beta 3$. *Cell.* 1996; 85: 683-93.
- 38) Morini M, Mottolèse M, Ferrari N, Ghiorzo F, Buglioni S, Mortarini R, et al. The $\alpha 3 \beta 1$ integrin is associated with mammary carcinoma cell metastasis, invasion, and gelatinase B (mmp - 9) activity. *Int J Cancer.* 2000; 87: 336-42.
- 39) Imai K, Hiramatsu A, Fukushima D, Pierschbacher MD, Okada Y. Degradation of decorin by matrix metalloproteinases: identification of the cleavage sites, kinetic analyses and transforming growth factor-beta1 release. *Biochem J.* 1997; 322: 809-14.
- 40) Mu D, Cambier S, Fjellbirkeland L, Baron JL, Munger JS, Kawakatsu H, et al. The integrin Mv8 mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF-t1. *J Cell Bio.* 2002; 157: 493-507.
- 41) Manicone AM, McGuire JK. Matrix Metalloproteinases as Modulators of Inflammation. *Semin Cell Dev Biol.* 2008; 19: 34-41.
- 42) Hiraoka N, Allen E, Apel IJ, Gyetko MR, Weiss SJ. Matrix Metalloproteinases Regulate

Neovascularization by Acting as Pericellular Fibrinolysins. *Cell*. 1998; 95: 365-77.

- 43) Mitsiades N, Yu WH, Poulaki V, Tsokos M, Stamenkovic I. Matrix metalloproteinase-7-mediated Cleavage of Fas Ligand Protects Tumor Cells From Chemotherapeutic Drug Cytotoxicity. *Cancer Res*. 2001; 61: 577-81.
- 44) Boudreau N, Sympson CJ, Werb Z, Bissell MJ. Suppression of ICE and Apoptosis in Mammary Epithelial Cells by Extracellular Matrix. *Science*. 1995; 267: 891-3.
- 45) Ferrari R, Martin G, Tagit O, Guichard A, Cambi A, Voituriez R, et al. MT1-MMP directs force-producing proteolytic contacts that drive tumor cell invasion. *Nat Commun*. 2019; 10: 4886.
- 46) Nowotny K, Grune T. Degradation of oxidized and glycoxidized collagen: role of collagen cross-linking. *Arch Biochem Biophys*. 2014; 542: 56-64.
- 47) Fosang AJ, Neame PJ, Last K, Hardingham TE, Murphy G, Hamilton JA. The interglobular domain of cartilage aggrecan is cleaved by PUMP, gelatinases, and cathepsin B. *J Biol Chem*. 1992; 267: 19470-19474.

■Chapter 9. Pathology and Disease

ITOH Yoshifumi

Kennedy Institute of Rheumatology, University of Oxford

- 1) Bella J, Hulmes DJ. Fibrillar Collagens. *Subcell Biochem.* 2017; 82: 457–90
- 2) Nagase H, Murphy G. Metalloproteinases in Cartilage Matrix Breakdown: The Roles in Rheumatoid Arthritis and Osteoarthritis. In: Brix K, Stöcker W, editors. *Proteases: Structure and Function.* Vienna: Springer Vienna; 2013. p. 433–69.
- 3) Ramirez F, Tanaka S, Bou-Gharios G. Transcriptional regulation of the human alpha2(I) collagen gene (COL1A2), an informative model system to study fibrotic diseases. *Matrix Biol.* 2006; 25: 365–72
- 4) Nagase H, Visse R. Triple Helicase Activity and the Structural Basis of Collagenolysis. *Extracellular Matrix Degradation* 2011. p. 95–122.
- 5) Ross FP, Christiano AM. Nothing but skin and bone. *J Clin Invest.* 2006; 116: 1140–9
- 6) Verzijl N, DeGroot J, Thorpe SR, Bank RA, Shaw JN, Lyons TJ, Bijlsma JW, Lafeber FP, Baynes JW, TeKoppele JM. Effect of collagen turnover on the accumulation of advanced glycation end products. *J Biol Chem.* 2000; 275: 39027–31
- 7) Liotta LA, Thorgeirsson UP, Garbisa S. Role of collagenases in tumor cell invasion. *Cancer Metastasis Rev.* 1982; 1: 277–88
- 8) Itoh Y, Nagase H. Matrix metalloproteinases in cancer. *Essays Biochem.* 2002; 38: 21–36
- 9) Itoh Y. Membrane-type matrix metalloproteinases: Their functions and regulations. *Matrix Biol.* 2015; 44–46: 207–23
- 10) Owen CA, Hu Z, Lopez-Otin C, Shapiro SD. Membrane-bound matrix metalloproteinase-8 on activated polymorphonuclear cells is a potent, tissue inhibitor of metalloproteinase-resistant collagenase and serpinase. *J Immunol.* 2004; 172: 7791–803
- 11) Sato H, Takino T, Okada Y, Cao J, Shinagawa A, Yamamoto E, Seiki M. A matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nature.* 1994; 370: 61–5
- 12) Itoh Y, Takamura A, Ito N, Maru Y, Sato H, Suenaga N, Aoki T, Seiki M. Homophilic complex formation of MT1-MMP facilitates proMMP-2 activation on the cell surface and promotes tumor cell invasion. *EMBO J.* 2001; 20: 4782–93
- 13) Holmbeck K, Bianco P, Caterina J, Yamada S, Kromer M, Kuznetsov SA, Mankani M, Robey PG, Poole AR, Pidoux I, Ward JM, Birkedal-Hansen H. MT1-MMP-deficient mice develop dwarfism, osteopenia, arthritis, and connective tissue disease due to inadequate collagen turnover. *Cell.* 1999; 99: 81–92
- 14) Holmbeck K, Bianco P, Yamada S, Birkedal-Hansen H. MT1-MMP: a tethered collagenase. *J Cell Physiol.* 2004; 200: 11–9

- 15) Novinec M, Lenarcic B. Cathepsin K: a unique collagenolytic cysteine peptidase. *Biol Chem.* 2013; 394: 1163–79
- 16) Garnero P, Borel O, Byrjalsen I, Ferreras M, Drake FH, McQueney MS, Foged NT, Delmas PD, Delaisse JM. The collagenolytic activity of cathepsin K is unique among mammalian proteinases. *J Biol Chem.* 1998; 273: 32347–52
- 17) Chung L, Dinakarpandian D, Yoshida N, Lauer-Fields JL, Fields GB, Visse R, Nagase H. Collagenase unwinds triple-helical collagen prior to peptide bond hydrolysis. *EMBO J.* 2004; 23: 3020–30
- 18) Manka SW, Carafoli F, Visse R, Bihan D, Raynal N, Farndale RW, Murphy G, Enghild JJ, Hohenester E, Nagase H. Structural insights into triple-helical collagen cleavage by matrix metalloproteinase 1. *Proc Natl Acad Sci U S A.* 2012; 109: 12461–6
- 19) Itoh Y, Ito N, Nagase H, Evans RD, Bird SA, Seiki M. Cell surface collagenolysis requires homodimerization of the membrane-bound collagenase MT1-MMP. *Mol Biol Cell.* 2006; 17: 5390–9
- 20) Itoh Y, Palmisano R, Anilkumar N, Nagase H, Miyawaki A, Seiki M. Dimerization of MT1-MMP during cellular invasion detected by fluorescence resonance energy transfer. *Biochem J.* 2011; 440: 319–26
- 21) Orgel JP, Irving TC, Miller A, Wess TJ. Microfibrillar structure of type I collagen in situ. *Proc Natl Acad Sci U S A.* 2006; 103: 9001–5
- 22) Sweeney SM, Orgel JP, Fertala A, McAuliffe JD, Turner KR, Di Lullo GA, Chen S, Antipova O, Perumal S, Ala-Kokko L, Forlino A, Cabral WA, Barnes AM, Marini JC, San Antonio JD. Candidate cell and matrix interaction domains on the collagen fibril, the predominant protein of vertebrates. *J Biol Chem.* 2008; 283: 21187–97
- 23) Perumal S, Antipova O, Orgel JP. Collagen fibril architecture, domain organization, and triple-helical conformation govern its proteolysis. *Proc Natl Acad Sci U S A.* 2008; 105: 2824–9
- 24) Zhu J, Hoop CL, Case DA, Baum J. Cryptic binding sites become accessible through surface reconstruction of the type I collagen fibril. *Sci Rep.* 2018; 8: 16646
- 25) Wynn TA. Fibrotic disease and the T(H)1/T(H)2 paradigm. *Nat Rev Immunol.* 2004; 4: 583–94
- 26) Theocharis AD, Manou D, Karamanos NK. The extracellular matrix as a multitasking player in disease. *FEBS J.* 2019; 286: 2830–69
- 27) Kinoshita T, Goto T. Molecular Mechanisms of Pulmonary Fibrogenesis and Its Progression to Lung Cancer: A Review. *Int J Mol Sci.* 2019; 20:
- 28) Tsochatzis EA, Bosch J, Burroughs AK. Liver cirrhosis. *Lancet.* 2014; 383: 1749–61
- 29) Denton CP, Khanna D. Systemic sclerosis. *Lancet.* 2017; 390: 1685–99
- 30) Lam WL, Rawlins JM, Karoo ROS, Naylor I, Sharpe DT. Re-Visiting Luck's Classification: A Histological Analysis of Dupuytren's Disease. *J Hand Surg-Eur Vol.* 2010; 35e: 312–7

- 31) Chandler C, Liu TS, Buckanovich R, Coffman LG. The double edge sword of fibrosis in cancer. *Transl Res.* 2019; 209: 55–67
- 32) Smolen JS, Aletaha D, McInnes IB. Rheumatoid arthritis. *Lancet.* 2016; 388: 2023–38
- 33) Nagase H, Murphy G. Metalloproteinases in cartilage Matrix Breakdown: The Roles in Rheumatoid Arthritis and Osteoarthritis. In: Brix K, Stöcker W, editors. *Proteases: Structure and Function*: Springer; 2014. p. 433–69.
- 34) Miller MC, Manning HB, Jain A, Troeberg L, Dudhia J, Essex D, Sandison A, Seiki M, Nanchahal J, Nagase H, Itoh Y. Membrane type 1 matrix metalloproteinase is a crucial promoter of synovial invasion in human rheumatoid arthritis. *Arthritis Rheum.* 2009; 60: 686–97
- 35) Majkowska I, Shitomi Y, Ito N, Gray NS, Itoh Y. Discoidin domain receptor 2 mediates collagen-induced activation of membrane-type 1 matrix metalloproteinase in human fibroblasts. *J Biol Chem.* 2017; 292: 6633–43
- 36) Glyn-Jones S, Palmer AJ, Agricola R, Price AJ, Vincent TL, Weinans H, Carr AJ. Osteoarthritis. *Lancet.* 2015; 386: 376–87
- 37) Karsdal, M. A., Madsen, S. H., Christiansen, C., Henriksen, K., Fosang, A. J. & Sondergaard, B. C. Cartilage degradation is fully reversible in the presence of aggrecanase but not matrix metalloproteinase activity. *Arthritis Res Ther* 10, R63, doi:10.1186/ar2434 (2008).
- 38) Xu L, Peng H, Wu D, Hu K, Goldring MB, Olsen BR, Li Y. Activation of the discoidin domain receptor 2 induces expression of matrix metalloproteinase 13 associated with osteoarthritis in mice. *J Biol Chem.* 2005; 280: 548–55
- 39) Compston JE, McClung MR, Leslie WD. Osteoporosis. *Lancet.* 2019; 393: 364–76
- 40) Al Anouti F, Taha Z, Shamim S, Khalaf K, Al Kaabi L, Alsafar H. An insight into the paradigms of osteoporosis: From genetics to biomechanics. *Bone Rep.* 2019; 11: 100216
- 41) Stetler-Stevenson WG, Aznavoorian S, Liotta LA. Tumor cell interactions with the extracellular matrix during invasion and metastasis. *Annu Rev Cell Biol.* 1993; 9: 541–73
- 42) Ohuchi E, Imai K, Fujii Y, Sato H, Seiki M, Okada Y. Membrane type 1 matrix metalloproteinase digests interstitial collagens and other extracellular matrix macromolecules. *J Biol Chem.* 1997; 272: 2446–51
- 43) Taniwaki K, Fukamachi H, Komori K, Ohtake Y, Nonaka T, Sakamoto T, Shiomi T, Okada Y, Itoh T, Itohara S, Seiki M, Yana I. Stroma-derived matrix metalloproteinase (MMP)-2 promotes membrane type 1-MMP-dependent tumor growth in mice. *Cancer Res.* 2007; 67: 4311–9
- 44) Hotary KB, Allen ED, Brooks PC, Datta NS, Long MW, Weiss SJ. Membrane type I matrix metalloproteinase usurps tumor growth control imposed by the three-dimensional extracellular matrix. *Cell.* 2003; 114: 33–45
- 45) Libby P, Buring JE, Badimon L, Hansson GK, Deanfield J, Bittencourt MS, Tokgozoglu L, Lewis EF. Atherosclerosis. *Nat Rev Dis Primers.* 2019; 5: 56

- 46) Schneider F, Sukhova GK, Aikawa M, Canner J, Gerdes N, Tang SM, Shi GP, Apte SS, Libby P. Matrix-metalloproteinase-14 deficiency in bone-marrow-derived cells promotes collagen accumulation in mouse atherosclerotic plaques. *Circulation*. 2008; 117: 931-9

■Chapter 10. Synthetic Collagen

KURODA Kazuki C., KOIDE Takaki

Department of Chemistry and Biochemistry, School of Advanced Science and Engineering, Waseda University

- 1) Koide T, Nishikawa Y, Takahara Y. Synthesis of heterotrimeric collagen models containing Arg residues in Y-positions and analysis of their conformational stability. *Bioorg Med Chem Lett.* 2004; 14: 125–8.
- 2) Holmgren SK, Taylor KM, Bretscher LE, Raines RT. Code for collagen's stability deciphered. *Nature.* 1998; 392: 666–7.
- 3) Russell LE, Fallas JA, Hartgerink JD. Selective assembly of a high stability AAB collagen heterotrimer. *J Am Chem Soc.* 2010; 132: 3242–3.
- 4) Jalan AA, Demeler B, Hartgerink JD. Hydroxyproline-free single composition ABC collagen heterotrimer. *J Am Chem Soc.* 2013; 135: 6014–7.
- 5) Xu F, Zhang L, Koder RL, Nanda V. De novo self-assembling collagen heterotrimers using explicit positive and negative design. *Biochemistry.* 2010; 49: 2307–16.
- 6) An B, Abbonante V, Xu H, Gavriilidou D, Yoshizumi A, Bihan D, et al. Recombinant collagen engineered to bind to discoidin domain receptor functions as a receptor inhibitor. *J Biol Chem.* 2016; 291: 4343–55.
- 7) Sakakibara S, Kishida Y, Kikuchi Y, Sakai R, Kakiuchi K. Synthesis of poly-(L-prolyl-L-prolyl-glycyl) of defined molecular weights. *B Chem Soc Jpn.* 1968; 41: 1273.
- 8) Sakakibara S, Inouye K, Shudo K, Kishida Y, Kobayashi Y, Prockop DJ. Synthesis of (Pro-Hyp-Gly)_n of defined molecular weights. Evidence for the stabilization of collagen triple helix by hydroxyproline. *Biochim Biophys Acta.* 1973; 303: 198–202.
- 9) Mechling DE, Bächinger HP. The collagen-like peptide (GER)15GPCCG forms pH-dependent covalently linked triple helical trimers. *J Biol Chem.* 2000; 275: 14532–6.
- 10) Shah NK, Ramshaw JAM, Kirkpatrick A, Shah C, Brodsky B. A host-guest set of triple-helical peptides: stability of Gly-X-Y triplets containing common nonpolar residues. *Biochemistry-US.* 1996; 35: 10262–8.
- 11) Persikov AV, Ramshaw JAM, Kirkpatrick A, Brodsky B. Amino acid propensities for the collagen triple-helix. *Biochemistry.* 2000; 39: 14960–7.
- 12) Persikov AV, Ramshaw JAM, Kirkpatrick A, Brodsky B. Peptide investigations of pairwise interactions in the collagen triple-helix. *J Mol Biol.* 2002; 316: 385–94.
- 13) Goodman M, Feng Y, Melacini G, Taulane JP. A template-induced incipient collagen-like triple-helical structure. *J Am Chem Soc.* 1996; 118: 5156–7.
- 14) Fields CG, Lovdahl CM, Miles AJ, Hageini VLM, Fields GB. Solid-Phase synthesis and stability

- of triple-helical peptides incorporating native collagen sequences. *Biopolymers*. 1993; 33: 1695-07.
- 15) Byrne C, McEwan PA, Emsley J, Fischer PM, Chan WC. End-stapled homo and hetero collagen triple helices: a click chemistry approach. *Chem. Commun.* 2011; 47: 2589-91.
 - 16) Ottl J, Moroder L. Disulfide-bridged heterotrimeric collagen peptides containing the collagenase cleavage site of collagen Type I. synthesis and conformational properties. *J Am Chem Soc.* 1999; 121: 653-61.
 - 17) Koide T, Yuguchi M, Kawakita M, Konno H. Metal-assisted stabilization and probing of collagenous triple helices. *J Am Chem Soc.* 2002; 124: 9388-9.
 - 18) Cai W, Kwok SW, Taulane JP, Goodman M. Metal-assisted assembly and stabilization of collagen-like triple helices. *J Am Chem Soc.* 2004; 126: 15030-1.
 - 19) Pires MM, Przybyla DE, Rubert Pérez CM, Chmielewski J. Metal-mediated tandem coassembly of collagen peptides into banded microstructures. *J Am Chem Soc.* 2011; 133: 14469-71.
 - 20) Pires MM, Lee J, Ernenwein D, Chmielewski J. Controlling the morphology of metal-promoted higher ordered assemblies of collagen peptides with varied core lengths. *Langmuir.* 2012; 28: 1993-7.
 - 21) Boudko S, Frank S, Kammerer RA, Stetefeld J, Schulthess T, Landwehr R, et al. Nucleation and propagation of the collagen triple helix in single-chain and trimerized peptides: transition from third to first order kinetics. *J Mol Biol.* 2002; 317: 459-70.
 - 22) Pakkanen O, Hämäläinen ER, Kivirikko KI, Myllyharju J. Assembly of stable human type I and III collagen molecules from hydroxylated recombinant chains in the yeast *Pichia pastoris*. Effect of an engineered C-terminal oligomerization domain foldon. *J Biol Chem.* 2003; 278: 32478-83.
 - 23) Jenkins CL, Bretscher LE, Guzei IA, Raines RT. Effect of 3-hydroxyproline residues on collagen stability. *J Am Chem Soc.* 2003; 125: 6422-7.
 - 24) Berisio R, Granata V, Vitagliano L, Zagari A. Imino acids and collagen triple helix stability: characterization of collagen-like polypeptides containing Hyp-Hyp-Gly sequence repeats. *J Am Chem Soc.* 2004; 126: 11402-3.
 - 25) Bann JG, Bächinger HP. Glycosylation/Hydroxylation-induced stabilization of the collagen triple helix. 4-trans-hydroxyproline in the Xaa position can stabilize the triple helix. *J Biol Chem.* 2000; 275: 24466-9.
 - 26) Persikov AV, Ramshaw JA, Brodsky B. Prediction of collagen stability from amino acid sequence. *J Biol Chem.* 2005; 280: 19343-9.
 - 27) Kramer RZ, Bella J, Mayville P, Brodsky B, Berman HM. Sequence dependent conformational variations of collagen triple-helical structure. *Nat Struct Biol.* 1999; 6: 454-7.
 - 28) Sekiya A, Okano-Kosugi H, Yamazaki CM, Koide T. Pigment epithelium-derived factor (PEDF)

- shares binding sites in collagen with heparin/heparan sulfate proteoglycans. *J Biol Chem.* 2011; 286: 26364–74.
- 29) Kawahara K, Yoshida T, Maruno T, Oki H, Ohkubo T, Koide T, Kobayashi Y. Spatiotemporal Regulation of PEDF Signaling by Type I Collagen Remodeling. *Proc Natl Acad Sci U S A* 2020; 117, 11450–11458.
- 30) Raynal N, Hamaia SW, Siljander PR, Maddox B, Peachey AR, Fernandez R, et al. Use of synthetic peptides to locate novel integrin $\alpha 2\beta 1$ -binding motifs in human collagen III. *J Biol Chem.* 2006; 281: 3821–31.
- 31) Farndale RW, Lisman T, Bihan D, Hamaia S, Smerling GS, Pugh N, et al. Cell-collagen interactions: the use of peptide Toolkits to investigate collagen-receptor interactions. *Biochem Soc Trans.* 2008; 36: 241–50.
- 32) Konitsiotis AD, Raynal N, Bihan D, Hohenester E, Farndale RW, Leitinger B. Characterization of high affinity binding motifs for the discoidin domain receptor DDR2 in collagen. *J Biol Chem.* 2008; 283: 6861–8.
- 33) Li Y, Foss CA, Summerfield DD, Doyle JJ, Torok CM, Dietz HC, et al. Targeting collagen strands by photo-triggered triple-helix hybridization. *Proc Natl Acad Sci USA.* 2012; 109: 14767–72.
- 34) Takita KK, Fujii KK, Kadonosono T, Masuda R, Koide T. Cyclic peptides for efficient detection of collagen. *Chembiochem.* 2018; 19: 1613–7.
- 35) Kar K, Amin P, Bryan MA, Persikov AV, Mohs A, Wang YH, et al. Self-association of collagen triple helix peptides into higher order structures. *J Biol Chem.* 2006; 281 :33283–90.
- 36) Xu F, Khan IJ, McGuinness K, Parmar AS, Silva T, Murthy NS, et al. Self-assembly of left- and right-handed molecular screws. *J Am Chem Soc.* 2013; 135: 18762–5.
- 37) Koide T, Homma DL, Asada S, Kitagawa K. Self-complementary peptides for the formation of collagen-like triple helical supramolecules. *Bioorg Med Chem Lett.* 2005; 15: 5230–3.
- 38) Kotch FW, Raines RT. Self-assembly of synthetic collagen triple helices. *Proc Natl Acad Sci USA.* 2006; 103: 3028–33.
- 39) O’Leary LER, Fallas JA, Bakota EL, Kang MK, Hartgerink JD. Multi-hierarchical self-assembly of a collagen mimetic peptide from triple helix to nanofibre and hydrogel. *Nat Chem.* 2011; 3: 821–8.
- 40) Kishimoto T, Morihara Y, Osanai M, Ogata S, Kamitakahara M, Ohtsuki C, et al. Synthesis of poly(Pro-Hyp-Gly) (n) by direct poly-condensation of (Pro-Hyp-Gly) (n), where n=1, 5, and 10, and stability of the triple-helical structure. *Biopolymers.* 2005; 79:163–72.
- 41) Suehiro T, Tada T, Waku T, Tanaka N, Hongo C, Yamamoto S, et al. Temperature-dependent higher order structures of the (Pro-Pro-Gly)₁₀-modified dendrimer. *Biopolymers.* 2011; 95: 270–7.
- 42) Ichise SF, Takeuchi S, Aoki S, Kuroda KC, Nose H, Masuda R, et al. Development of a collagen-like peptide polymer via end-to-end disulfide cross-linking and its application as a biomaterial.

Acta Biomater. 2019; 94: 361–71.

- 43) Luo J, Tong YW. Self-assembly of collagen-mimetic peptide amphiphiles into biofunctional nanofiber. *ACS Nano*. 2011; 5: 7739–47
- 44) Krishna OD, Jha AK, Jia X, Kiick KL. Integrin-mediated adhesion and proliferation of human MSCs elicited by a hydroxyproline-lacking, collagen-like peptide. *Biomaterials*. 2011; 32: 6412–24.
- 45) Luo T, Kiick KL. Noncovalent modulation of the inverse temperature transition and self-assembly of elastin-b-collagen-like peptide bioconjugates. *J Am Chem Soc*. 2015; 137: 15362–5.
- 46) Stahl PJ, Romano NH, Wirtz D, Yu SM. PEG-based hydrogels with collagen mimetic peptide-mediated and tunable physical cross-links. *Biomacromolecules*. 2010; 11: 2336–44
- 47) Rubert Pérez CM, Rank LA, Chmielewski J. Tuning the thermosensitive properties of hybrid collagen peptide-polymer hydrogels. *Chem Commun (Camb)*. 2014; 50: 8174–6.
- 48) Luo T, David MA, Dunshee LC, Scott RA, Urello MA, Price C, et al. Thermoresponsive elastin-b-collagen-like peptide bioconjugate nanovesicles for targeted drug delivery to collagen-containing matrices. *Biomacromolecules*. 2017; 18: 2539–51.
- 49) Kumar VA, Taylor NL, Jalan AA, Hwang LK, Wang BK, Hartgerink JD. A nanostructured synthetic collagen mimic for hemostasis. *Biomacromolecules*. 2014; 15: 1484–90.

■Chapter 11. Digestion and Absorption

SATO Kenji

Marine Biological Function, Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University

SHIGEMURA Yasutaka

Department of domestic science, Tokyo Kasei University

- 1) Sato K, Yoshinaka R, Sato M, Shimizu Y. Isolation of native acid-soluble collagen from fish muscle. *Nippon Suisan Gakkaishi*. 1987; 53: 1431-1436.
- 2) Rudenskaya GN, Isaev VA, Stepanov VM, Dunaevsky YE, Baratova LA, Kalebina TS, et al. Isolation and properties of serine proteinase PC of the kamchatka crab *Paralithodes camchatica*, a proteolytic enzyme of broad specificity. *Biochemistry*. 1996; 61: 804-814.
- 3) Robinson JJ. Proteolytic processing of a sea urchin, ECM-localized protein into lower mol mass species possessing collagen-cleavage activity. *J Cell Biochem*. 2006; 99: 816-823.
- 4) 菊地数晃, 又平芳春. ヒトの乾燥肌および肌荒れに対する海洋性コラーゲンペプチド含有飲料の有用性. *Fragrance Journal*. 2003; 31: 97-102.
- 5) 福岡伸一. 動的平衡: 生命はなぜそこに宿るのか. 木楽舎: 2009
- 6) Iwai K, Hasegawa T, Taguchi Y, Morimatsu F, Sato K, Nakamura Y, et al. Identification of food-derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. *J Agric Food Chem*. 2005; 53: 6531-6536.
- 7) Ohara H, Matsumoto H, Ito K, Iwai K, Sato K. Comparison of quantity and structures of hydroxyproline-containing peptides in human blood after oral ingestion of gelatin hydrolysates from different sources. *J. Agric. Food Chem*. 2007; 55: 1532-1535.
- 8) Shigemura Y, Kubomura D, Sato Y, Sato K. Dose-dependent changes in the levels of free and peptide forms of hydroxyproline in human plasma after collagen hydrolysate ingestion. *Food Chem*. 2014; 159: 328-332.
- 9) 佐藤健司, 依田悠里. ペプチドの吸収 宮澤陽夫監修 食品機能性成分の吸収・代謝機構, シーエムシー出版: 2013
- 10) 市川聡美, 大原浩樹, 大庭知慧, 伊藤恭子, 松本均, 竹内康雄ら. 平均分子量の異なるコラーゲンペプチド経口摂取がペプチド型ヒドロキシプロリンの血中移行量に及ぼす影響. *医学と薬学*. 2009; 62: 801-807.
- 11) Taga Y, Iwasaki Y, Shigemura Y, Mizuno K. Improved in vivo tracking of orally administered collagen hydrolysate using stable isotope labeling and LC-MS techniques. *J Agric Food Chem*. 2019; 67: 4671-4678.
- 12) Shigemura Y, Akaba S, Kawashima E, Park EY, Nakamura Y, Sato K. Identification of a novel

- food-derived collagen peptide, hydroxyprolyl-glycine, in human peripheral blood by pre-column derivatisation with phenyl isothiocyanate. *Food Chem.* 2011; 129: 1019–1024.
- 13) Shigemura Y, Suzuki A, Kurokawa M, Sato Y, Sato K. Changes in composition and content of food-derived peptide in human blood after daily ingestion of collagen hydrolysate for 4 weeks. *J Sci Food Agric.* 2018; 98: 1944–1950.
 - 14) Aito-Inoue M, Lackeyram D, Fan MZ, Sato K, Mine Y. Transport of a tripeptide, Gly-Pro-Hyp, across the porcine intestinal brush-border membrane. *J Pept Sci.* 2007; 13: 468–474.
 - 15) Shimizu K, Sato M, Zhang Y, Kouguchi T, Takahata Y, Morimatsu F, et al. The bioavailable octapeptide Gly-Ala-Hyp-Gly-Leu-Hyp-Gly-Pro stimulates nitric oxide synthesis in vascular endothelial cells. *J Agric Food Chem.* 2010; 58: 6960–6965.
 - 16) Absorption and metabolism of orally administered collagen hydrolysates evaluated by the vascularly perfused rat intestine and liver in situ. *Biomed Res.* 2018; 39(1): 1–11.
 - 17) Taga Y, Kusubata M, Ogawa-Goto K, Hattori S. Efficient absorption of X-hydroxyproline (Hyp)-Gly after oral administration of a novel gelatin hydrolysate prepared using ginger protease. *J Agric Food Chem.* 2016; 64: 2962–2970.
 - 18) Asai T, Takahashi A, Ito K, Uetake T, Matsumura Y, Ikeda K, et al. Amount of collagen in the meat contained in Japanese daily dishes and the collagen peptide content in human blood after ingestion of cooked fish meat. *J Agric Food Chem.* 2019; 67: 2831–2838.
 - 19) Kusubata M, Koyama Y, Tometsuka C, Shigemura Y, Sato K. Detection of endogenous and food-derived collagen dipeptide prolylhydroxyproline (Pro-Hyp) in allergic contact dermatitis-affected mouse ear. *Biosci Biotechnol Biochem.* 2015; 79: 1356–1361.
 - 20) Asai T, Yoshikawa K, Sawada K, Fukamizu K, Koyama Y, Shigemura Y, et al. Mouse skin fibroblasts with mesenchymal stem cell marker p75 neurotrophin receptor proliferate in response to prolyl-hydroxyproline. *J Func Foods.* 2020; 66: 103792.
 - 21) Abe H, Satoh M, Miyauchi S, Shuto, S, Matsuda, A, Kamo, N. Conjugation of dipeptide to fluorescent dyes enhances its affinity for a dipeptide transporter (PEPT1) in human intestinal Caco-2 cells. *Bioconjugate Chem.* 1999; 10: 24–31.
 - 22) Kawaguchi T, Nanbu PN, Kurokawa M. Distribution of prolylhydroxyproline and its metabolites after oral administration in rats. *Biol Pharm Bull.* 2012; 35: 422–427.
 - 23) Kimira Y, Odaira H, Nomura K, Taniuchi Y, Inoue N, Nakatani S, et al. Collagen-derived dipeptide prolyl-hydroxyproline promotes osteogenic differentiation through Foxg1. *Cell Mol Biol Lett.* 2017; 22: 27–38.
 - 24) Nakatani S, Mano H, Sampei C, Shimizu J, Wada M. Chondroprotective effect of the bioactive peptide prolyl-hydroxyproline in mouse articular cartilage in vitro and in vivo. *Osteoarthritis Cartilage.* 2009; 17: 1620e1627.
 - 25) Ohara H, Iida H, Ito K, Takeuchi Y, Nomura Y. Effects of Pro-Hyp, a collagen hydrolysate-

derived peptide, on hyaluronic acid synthesis using in vitro cultured synovium cells and oral ingestion of collagen hydrolysates in a guinea pig model of osteoarthritis. *Biosci Biotechnol Biochem.* 2010; 74: 2096–2099.

- 26) Shigemura Y, Iwai K, Morimatsu F, Iwamoto T, Mori T, Oda T, et al. Effect of Prolyl-hydroxyproline (Pro-Hyp), a food-derived collagen peptide in human blood, on growth of fibroblasts from mouse skin. *J Agric Food Chem.* 2009; 57: 444–449.

■Chapter 12. Mammalian collagen

TANAKA Keisuke, HATTORI Shunji

Nippi Research Institute of Biomatrix

References

- (1) Ricard-Blum S. The Collagen Family. *Cold Spring Harb Perspect Biol.* 2011; 3(1): a004978.
- (2) Kumazawa Y, Taga Y, Iwai K, Koyama Y. A Rapid and Simple LC-MS Method Using Collagen Marker Peptides for Identification of the Animal Source of Leather. *J. Agric. Food Chem.* 2016; 64: 6051-7.
- (3) Ebihara T, Iijima K, Sato K, Someki I, Kuwaba K, Hattori S, Irie S. Changes in the Extractability of Bovine Skin Collagen through the Course of Aging and Biochemical Properties of the Collagen Extracted (in Japanese). *Connective Tissue.* 1999; 31: 17-23.
- (4) Hattori S, Adachi E, Ebihara T, Shirai T, Someki I, Irie S. Alkali-treated collagen retained the triple helical conformation and the ligand activity for the cell adhesion via alpha2beta1 integrin. *J Biochem.* 1999; 125(4): 676-84.
- (5) Ushiki Y, Hattori S, Irie S. Safety aspect of collagen as a raw material for cosmetics (in Japanese). *Fragrance Journal.* 1997; 25(7): 42-47.
- (6) Muraoka M, Nakazato K, Hayashi T. Gel formation from the collagen type IV isolated from bovine lens capsule in guanidine-HCl and dithiothreitol. *J Biochem.* 1996; 119(1): 167-72.
- (7) Burjanadze TV. Evidence for the role of 4-hydroxyproline localized in the third position of the triplet (Gly-X-Y) in adaptational changes of thermostability of a collagen molecule and collagen fibrils. *Biopolymers.* 1982; 21: 1489-1501.
- (8) Sato K, Ebihara T, Adachi E, Kawashima S, Hattori S, Irie S. Possible involvement of aminotelopeptide in self-assembly and thermal stability of collagen I as revealed by its removal with proteases. *J Biol Chem.* 2000; 275(33): 25870-5.
- (9) Yamauchi M, Sricholpech M. Lysine post-translational modifications of collagen. *Essays Biochem.* 2012; 52: 113-33.
- (10) Taga Y, Kusubata M, Ogawa-Goto K, Hattori S. Stable isotope-labeled collagen: a novel and versatile tool for quantitative collagen analyses using mass spectrometry. *J Proteome Res.* 2014; 13(8): 3671-8.
- (11) Koyama Y. Functional aspects of collagen. *Hikakukagaku* 2010; 56(2): 71-9.
- (12) Zhang Z, Liu W, Li D, Li G. Physicochemical properties of succinylated calfskin pepsin-solubilized collagen. *Biosci Biotechnol Biochem.* 2007; 71(8): 2057-60.
- (13) Hattori S, Adachi E, Ebihara T, Shirai T, Someki I, Irie S. Alkali-treated collagen retained the triple helical conformation and the ligand activity for the cell adhesion via alpha2beta1 integrin. *J Biochem.* 1999; 125(4): 676-84.

- (14) Miyata T, Taira T, Noishiki, Y. Collagen engineering for biomaterial use. *Clin Mater.* 1992; 9: 139–48
- (15) Stenzel KH, Dunn MW, Rubin AL, Miyata T. Collagen gels: design for a vitreous replacement. *Science.* 1969; 164(3885): 1282–3.
- (16) Mimura T, Yamagami S, Yokoo S, Usui T, Tanaka K, Hattori S, et al. Cultured human corneal endothelial cell transplantation with a collagen sheet in a rabbit model. *Invest Ophthalmol Vis Sci.* 2004; 45(9): 2992–7.
- (17) Noritake K, Kuroda S, Nyan M, Atsuzawa Y, Uo M, Ohya K, et al. Use of a gelatin hydrogel membrane containing β -tricalcium phosphate for guided bone regeneration enhances rapid bone formation. *Dent Mater J.* 2014; 33(5): 674–80.
- (18) Lee A, Hudson AR, Shiwarski DJ, Tashman JW, Hinton TJ, Yerneni S, et al. 3D bioprinting of collagen to rebuild components of the human heart. *Science.* 2019; 365(6452): 482–7.
- (19) Tanaka K, Ogura T, Hattori S, Matsuda K, Kobayashi Y, Oi S. Collagen powder and/or collagen-derived powder, and production method for the same. US Patent No.: US 9,617,298, Date of patent: 2017.4.11
- (20) Ogura T, Tanaka K, Ohba Y, Hattori S. Collagen structure, and method for producing collagen structure. US application pub. No.: US 2015/0004414 Pub. Date: 2015.1.1
- (21) Uchida K, Matsushita O, Naruse K, Mima T, Nishi N, Hattori S, et al. Acceleration of periosteal bone formation by human basic fibroblast growth factor containing a collagen-binding domain from *Clostridium histolyticum* collagenase. *J Biomed Mater Res A.* 2014; 102(6): 1737–43.
- (22) Saito W, Uchida K, Ueno M, Matsushita O, Inoue G, Nishi N, et al. Acceleration of bone formation during fracture healing by injectable collagen powder and human basic fibroblast growth factor containing a collagen-binding domain from *Clostridium histolyticum* collagenase. *J Biomed Mater Res A.* 2014; 102(9): 3049–55
- (23) Nakamura S, Ito T, Okamoto K, Mima T, Uchida K, Siddiqui YD, et al. Acceleration of bone regeneration of horizontal bone defect in rats using collagen-binding basic fibroblast growth factor combined with collagen scaffolds. *J Periodontol.* 2019; 90(9): 1043–1052.

■Chapter 13. Collagen from Aquatic Animals

ISHIHARA Kenji

Fisheries Technology Institute, Japan Fisheries Research and Education Agency

HIRAOKA Yoshinobu

Ehime Institute of Invention

- 1) Ikoma T, Kobayashi H, Tanaka J, Walsh D, Mann S. Physical properties of type I collagen extracted from fish scales of *Pagrus major* and *Oreochromis niloticus*. *Int J Biol Macromol*. 2003; 32: 199-204.
- 2) 服部俊治, 動物由来繊維分子コラーゲンの性質と応用, 繊維と工業. 2009; 65: 453-461.
- 3) 西塔正孝, I-3. 魚類のコラーゲン. 日本水産学会誌. 2017; 83: 818.
- 4) 木村茂, 第2章 脊索動物(魚類)-酸可溶性コラーゲンの宝庫-. 「海洋性コラーゲンを探る」, 五曜出版(東京). 2014; pp. 29-46.
- 5) 高橋豊雄・田中武夫, 水産皮革の理化学的研究-Ⅷ. 魚皮革の耐熱性に就いて 其の三). 日本水産学会誌. 1953; 19: 603-609.
- 6) Saito M, Takenouchi Y, Kunisaki N, Kimura S. Complete primary structure of rainbow trout type I collagen consisting of 1(I)12(I)23(I) heterotrimers. *Eur. J. Biochem*. 2001; 268: 2817-2827.
- 7) 高橋豊雄, 横山和吉, 水産皮革の理化学的研究-Ⅶ. 魚皮コラーゲンのヒドロキシプロリン含量. 日本水産学会誌. 1954; 20: 525-529.
- 8) 畑江 敬子, 飛松 聡子, 竹山 まゆみ, 松本 重一郎, 魚肉の物性とその魚種差に対する結合組織の寄与. 日本水産学会誌. 1986; 52: 2001-2007.
- 9) 佐藤健司・坂口守彦, 魚介類のおいしさの秘密. 化学と生物. 1998; 36: 504-509.
- 10) 畑江敬子, 9. 調理と細胞外マトリックス. 「魚介類の細胞外マトリックス」, 木村茂編, 恒星社厚生閣(東京). 1997; pp. 91-101.
- 11) 木村茂, 第5章 棘皮動物-ヒトデ, ウニ, ナマコ-. 「海洋性コラーゲンを探る」, 五曜出版(東京). 2014; pp. 75-92.
- 12) 木村茂, 第6章 節足動物-カニ, エビ-. 「海洋性コラーゲンを探る」, 五曜出版(東京). 2014; pp. 93-100.
- 13) 木村茂, 第8章 軟体動物-タコ, イカ, アワビなど-. 「海洋性コラーゲンを探る」, 五曜出版(東京). 2014; pp. 123-138.
- 14) 木村茂, 第9章 刺胞動物-エチゼンクラゲ-. 「海洋性コラーゲンを探る」, 五曜出版(東京). 2014; pp. 139-156.
- 15) <http://www.fra.affrc.go.jp/pressrelease/pr18/190316/kurage4.pdf>
- 16) https://www.gmj.or.jp/collagen_factbook/HTML/list1.html
- 17) 健康食品の市場動向と素材・技術研究. 食品と開発. 2010; 45: 25-48.

- 18) 吉仲禮二, 8. 魚類のコラーゲン, 「水産動物筋肉タンパク質の比較生化学」、新井健一 編 恒星社厚生閣(東京). 1989; pp. 81-99.
- 19) 石原賢司・小山田千秋・土居幹治・松本淳一・白石隆介・平岡芳信・園田浩二, カツオの中骨を原料とするコラーゲンペプチドの製造方法. 特開 2012-244930.
- 20) 雑賀愛・岩井浩二・高畑能久, 血圧上昇抑制作用を有するペプチド. 特願 2008-506354.
- 21) Iwai K, Hasegawa T, Taguchi Y, Morimatsu F, Sato K, Nakamura Y, Higashi A, Kido Y, Nakabo Y, Ohtsuki K, Identification of food-derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. J. Agric. Food Chem. 2005; 53: 6531-6536.
- 22) 重村泰毅, 魚類由来タンパク質加水分解物摂取後のヒト体調改善効果の予測されるメカニズム. 日本食生活学会誌. 2016; 27: 147-152.

■Chapter 14. Collagen from lower animals

ARAI Katsuhiko

Scleroprotein and Leather Research Institute, Faculty of Agriculture, Tokyo University of Agriculture and Technology

- 1) Parisi JR, Fernandes KR, Avanzi IR, Dorileo BP, Santana AF, Andrade AL, Gabbai-Armelin PR, Fortulan CA, Trichês ES, Granito RN, Renno ACM. Incorporation of Collagen from Marine Sponges (Spongin) into Hydroxyapatite Samples: Characterization and In Vitro Biological Evaluation. *Mar Biotechnol (NY)*. 2019. 21: 30–37.
- 2) Draper GW, Shoemark DK, Adams JC. Modelling the early evolution of extracellular matrix from modern Ctenophores and Sponges. *Essays Biochem*. 2019. 63:389–405.
- 3) Boute N, Exposito JY, Boury-Esnault N, Vacelet J, Noro N, Miyazaki K, Yoshizato K, Garrone R. Type IV collagen in sponges, the missing link in basement membrane ubiquity. *Biol Cell*. 1996. 88:37–44.
- 4) Exposito JY, Cluzel C, Garrone R, Lethias C. Evolution of collagens. *Anat Rec*. 2002. 268: 302–316.
- 5) Aouacheria A, Geourjon C, Aghajari N, Navratil V, Deléage G, Lethias C, Exposito JY. Insights into early extracellular matrix evolution: spongin short chain collagen-related proteins are homologous to basement membrane type IV collagens and form a novel family widely distributed in invertebrates. *Mol Biol Evol*. 2006. 23: 2288–2302.
- 6) Ehrlich H, Wysokowski M, Żółtowska-Aksamitowska S, Petrenko I, Jesionowski T. Collagens of Poriferan Origin. *Mar Drugs*. 2018. 16: E79.
- 7) Tziveleka LA, Ioannou E, Tsiourvas D, Berillis P, Foufa E, Roussis V. Collagen from the Marine Sponges *Axinella cannabina* and *Suberites carnosus*: Isolation and Morphological, Biochemical, and Biophysical Characterization. *Mar Drugs*. 2017. 15: E152.
- 8) Miura S, Kimura S. Jellyfish mesogloea collagen. Characterization of molecules as alpha 1 alpha 2 alpha 3 heterotrimers. *J Biol Chem*. 1985. 260:15352–15356.
- 9) Addad S, Exposito JY, Faye C, Ricard-Blum S, Lethias C. Isolation, characterization and biological evaluation of jellyfish collagen for use in biomedical applications. *Mar Drugs*. 2011. 9: 967–983.
- 10) Miki A, Inaba S, Baba T, Kihira K, Fukada H, Oda M. Structural and physical properties of collagen extracted from moon jellyfish under neutral pH conditions. *Biosci Biotechnol Biochem*. 2015. 79: 1603–1607.
- 11) Wang W, Omori M, Hayashibara T, Shimoike K, Hatta M, Sugiyama T, Fujisawa T. Isolation and characterization of a mini-collagen gene encoding a nematocyst capsule protein from a reef-building coral, *Acropora donei*. *Gene*. 1995. 152: 195–200.

- 12) Engel U, Pertz O, Fauser C, Engel J, David CN, Holstein TW. A switch in disulfide linkage during minicollagen assembly in *Hydra* nematocysts. *EMBO J.* 2001. 20: 3063–3073.
- 13) Shpirer E, Chang ES, Diamant A, Rubinstein N, Cartwright P1, Huchon D. Diversity and evolution of myxozoan minicollagens and nematogalectins. *BMC Evol Biol.* 2014. 14:205.
- 14) Yoneda C, Ahsan MN, Nakaya M, Matsubara Y, Ebihara T, Irie S, Uno Y, Hatae K, Watabe S. Abalone collagens: immunological properties and seasonal changes of their mRNA levels. *Comp Biochem Physiol B Biochem Mol Biol.* 2000. 126: 59–68.
- 15) Badiu DL, Luque R, Dumitrescu E, Craciun A, Dinca D. Amino acids from *Mytilus galloprovincialis* (L.) and *Rapana venosa* molluscs accelerate skin wounds healing via enhancement of dermal and epidermal neoformation. *Protein J.* 2010 29:81–92.
- 16) Kimura S, Nagaoka Y, Kubota M. Studies on marine invertebrate collagens- I. Some collagens from crustaceans and molluscs. *Bull. Japan. Soc. Sci. Fish.* 1969. 35: 743–748.
- 17) Nagai T. Collagen from diamondback squid (*Thysanoteuthis rhombus*) outer skin. *Z Naturforsch C.* 2004. 59:271–275.
- 18) Sivakumar P, Chandrakasan G. Occurrence of a novel collagen with three distinct chains in the cranial cartilage of the squid *Sepia officinalis*: comparison with shark cartilage collagen. *Biochim Biophys Acta.* 1998. 1381:161–169.
- 19) Kimura S, Takema Y, Kubota M. Octopus skin collagen. Isolation and characterization of collagen comprising two distinct alpha chains. *J Biol Chem.* 1981. 256:13230–13234.
- 20) Johnstone IL. Cuticle collagen genes. Expression in *Caenorhabditis elegans*. *Trends Genet.* 2000. 16:21–27.
- 21) Ashhurst DE, Bailey AJ. Locust collagen: morphological and biochemical characterization. *Eur J Biochem.* 1980. 103: 75–83.
- 22) Cecchini JP, Knibiehler B, Mirre C, Le Parco Y. Evidence for a type-IV-related collagen in *Drosophila melanogaster*. Evolutionary constancy of the carboxyl-terminal noncollagenous domain. *Eur J Biochem.* 1987. 165: 587–593.
- 23) Wang X, Harris RE, Bayston LJ, Ashe HL. Type IV collagens regulate BMP signalling in *Drosophila*. *Nature.* 2008. 455: 72–77.
- 24) Baerwald RJ, Williamson LC, Stevens E, Rike C, Trabanino S, Carlton J. Isolation, ultrastructure, and partial characterization of collagen from the perineurium of the Florida lobster, *Panulirus argus*. *Biochem Cell Biol.* 1991. 69: 531–536.
- 25) Sivakumar P, Suguna L, Chandrakasan G. Molecular species of collagen in the intramuscular connective tissues of the marine crab, *Scylla serrata*. *Comp Biochem Physiol B Biochem Mol Biol.* 2000. 125:555–562.
- 26) Trotter JA, Thurmond FA, Koob TJ. Molecular structure and functional morphology of echinoderm collagen fibrils. *Cell Tissue Res.* 1994. 275: 451–458.

- 27) Motokawa T. Connective tissue catch in echinoderms. *Biol Rev* 1984. 59: 255–270.
- 28) Adibzadeh N, Aminzadeh S, Jamili S, Karkhane AA, Farrokhi N. Purification and characterization of pepsin-solubilized collagen from skin of sea cucumber *Holothuria parva*. *Appl Biochem Biotechnol*. 2014. 173: 143–154.
- 29) Zhong M, Chen T, Hu C, Ren C. Isolation and characterization of collagen from the body wall of sea cucumber *Stichopus monotuberculatus*. *J Food Sci*. 2015. 80:C671–679.
- 30) Ovaska M, Bertalan Z, Miksic A, Sugni M, Di Benedetto C, Ferrario C, Leggio L, Guidetti L, Alava MJ, La Porta CAM, Zapperi S. Deformation and fracture of echinoderm collagen networks. *J Mech Behav Biomed Mater*. 2017. 65: 42–52.

■Chapter 15. Application of Collagen, Overview

NOMURA Yoshihiro

Scleroprotein and Leather Research Institute, Faculty of Agriculture, Tokyo University of Agriculture and Technology

- 1) 野村義宏 コラーゲン今昔物語～はじめに～ 皮革科学 58(2) 47-50(2012)
- 2) 竹ノ内一昭 古代の皮革 4. 東・南アジア 皮革科学 1-5(200)
- 3) 竹ノ内一昭 古代の皮革 5 東・日本 皮革科学 1-6(200)
- 4) 千賀久 古墳時代の革製品 皮革科学 50, 129 (2004)
- 5) Egypt unveils biggest ancient coffin find in over a century, REUTERS Oct 19, 2019
<https://www.reuters.com/article/us-egypt-archaeology-luxor/egypt-unveils-biggest-ancient-coffin-find-in-over-a-century-idUSKBN1WY0C5>
- 6) 綿谷正之 墨の文化史 概説 奈良保育学院 研究紀要 1-34 (2014)
- 7) 高田俊二 写真産業の技術革新史—その 1 乳剤時代の幕開けと感光材料の工業化 日本写真学会誌 77(2), 76-82(2014)
- 8) RL Madox, British J Photo, 18 422(1871)
- 9) 野村義宏 コラーゲン今昔物語～コラーゲンの利用分野～ 皮革科学 58(3) 93-99(2012)
- 10) T Nishihara, T Miyata, Collagen symposium, 3, 66-93 (1962)
- 11) T Fujii, Hoppe-Seylers Z Physiol Chem, 350, 1257-1265 (1969)
- 12) 野村義宏 コラーゲン今昔物語～医療用途でのコラーゲンの活用～ 皮革科学 59(3), 99-106 (2013)
- 13) 野村義宏 コラーゲン今昔物語～可食性フィルム～ 皮革科学 60(1) 1-5 (2014)
- 14) E Bell, B Ivarsson, C Merrill, Proc Natl Acad Sci USA, 76, 1274-1278 (1979)

■Chapter 16. Leather and Its Products

YOSHIMURA Keiji

Japan Leather and Leather Goods Industries Association

- 1) O'Sullivan NJ, Teasdale MD, Mattiangeli V, Maixner F, Pinhasi R, Bradley DG, Zink A. A whole mitochondria analysis of the Tyrolean Iceman's leather provides insights into the animal sources of Copper Age clothing. *Scientific Reports*. 2016; 6, 31279
- 2) <https://leathercouncil.org/information/statistics-sources-of-information/>
- 3) www.fao.org/faostat/ www.fao.org/faostat/
- 4) 日本皮革技術協会皮革ハンドブック編集委員会, 皮革ハンドブック, 樹芸書房, 2005, p24.
- 5) Buljan J, Kral I. The framework for sustainable leather manufacture. 2nd ed. UNIDO, 2018. p. 22.
- 6) Covington AD. Tanning chemistry, The science of leather, The Royal Society of Chemistry (UK); 2009, p284.
- 7) 日本皮革技術協会, 新版皮革科学, 日本皮革技術協会, 1992, p53-56.
- 8) <http://japan-ecoleather.jp/%e6%97%a5%e6%9c%ac%e3%82%a8%e3%82%b3%e3%83%ac%e3%82%b6%e3%83%bc%e3%81%a8%e3%81%af/%ef%bc%96%e3%81%a4%e3%81%ae%e6%9d%a1%e4%bb%b6/>
- 9) Ogata K, Kumazawa K, Koyama Y, Yoshimura K, Takahashi K, Complete inhibition of hexavalent chromium formation from chrome-tanned leather with combined inhibitors, *J Soc Leather Technol Chem*, 101, 27-32, 2017.
- 10) Ogata K, Kumazawa K, Hattori S, Yoshimura K, Takahashi K, Self-conversion of hexavalent chromium formed in chrome-tanned leather during long-term storage and perfect inhibition with a combination of inhibitors, *J Soc Leather Technol Chem*, 102, 53-58, 2018.
- 11) Ogata K, Inatsugi Y, Hattori S, Kumazawa K, Yoshimura K, Takahashi K, Pilot production of chrome-tanned leather without formation of hexavalent chromium by treatment with a combination of inhibitors, *J Soc Leather Technol Chem*, 103, 1-6, 2019.

■Chapter 17. Japanese glue and Western glue

DOI Masahiro

Asahi Gelatine Industrial Co., Ltd.

- 1) <https://dictionary.jlia.or.jp/detail.php?id=1933>
- 2) <http://urakami.com/nikawa.html>
- 3) パギイ法 写真用ゼラチン試験法 第10版(2006年度版)
- 4) 日本にかわ・ゼラチン工業組合：にかわとゼラチン
- 5) 膠文化財研究会：膠の基礎知識
- 6) 渡部睦人、野村義宏：カラーゲン今昔物語 膠とゼラチン
- 7) 関出：日本画とにかわ 兵庫県バイオポリマー研究会
- 8) 土居昌裕：三千本膠「飛鳥」の開発について 愛知県立大学講演要旨
- 9) 宇高健太郎, 野村義宏, 寺岡葉子：膠製造における諸条件と製品の性状の関連(3) -製造条件と物性決定因子- 文化財保存修復学会第34回大会研究発表要旨集; 2012
- 10) 大野隆司：写真用ゼラチンの最近の傾向 日本写真学会誌 52, 320-322 (1989)
- 11) 伊藤典一：最近の写真用ゼラチンの製法 日本写真学会誌 52, 329-331 (1989)
- 12) 戸田雄三：銀塩感光材料における写真用ゼラチンの役割 日本写真学会誌 52, 323-328(1989)
- 13) 久米裕二：国立博物館 技術の系統化調査報告書 第17集、カラーネガフィルムの技術系統化調査 国立科学博物館 産業技術史資料情報センター 275-363 (2012)
- 14) 高田俊二：写真産業の技術革新史-その1 乳剤時代の幕開けと感光材料の工業化 日本写真学会誌 77, 76-82(2014)
- 15) 高田俊二：写真産業の技術革新史-その2 乳剤高感度化技術と写真・映像の大衆化 日本写真学会誌 78, 23-31(2015)
- 16) 小林 隆：ゼラチンの物性評価とその標準化 日本写真学会誌 67, 386-391 (2004)

■Chapter 18. Inkstick (sumi) and Classical Animal Glue

UDAKA Kentaro

Independent Administrative Institution National Institutes for Cultural Heritage, Tokyo National Research Institute for Cultural Properties

- 1) TERADA Torahiko, YAMAMOTO Ryuzo. Experimental studies on colloid nature of Chinese black ink Part I. Scientific Papers of the Institute of Physical and Chemical Research 23; 1934
- 2) TERADA Torahiko, YAMAMOTO Ryuzo. Experimental Studies on Colloid Nature of Chinese Black Ink, Part II, Scientific Papers of the Institute of Physical and Chemical Research 27; 1935
- 3) 飯島茂. 硯墨新語. 雄山閣; 1935
- 4) 宮坂和雄. 墨と色材の知識. 木耳社; 1972
- 5) 宮坂和雄, 齊藤昌子, 有馬智加子, 片岡友子. 墨の粒子の定着状態と墨色に関する研究. 共立女子大学家政学部紀要 20; 1974
- 6) 小口八郎. 墨の研究. 古文化財之科学 文化財保存修復学会誌 20・21 (1977)
- 7) 編 宇野健一. 徳川美術館蔵 古墨. しこうしゃ図書販売; 1991
- 8) Joseph R. Swider, Vincent A. Hackley, John Winter. Characterization of Chinese Ink in Size and Surface. Journal of Cultural Heritage 4; 2003
- 9) 日野楠雄. 墨色の変化 -紙と墨と水と- 宣紙～楮紙へ、にじみの世界. 和紙文化研究 17; 2009
- 10) 為近磨巨登. 写真で知る墨・硯・紙. 木耳社; 2009
- 11) 為近磨巨登. 書道用紙とにじみ. 木耳社; 2011
- 12) 宇高健太郎. 古典絵画における墨の研究 付論：伊藤若冲紙本墨画作品の復元模写による検証. 東京藝術大学博士研究論文; 2013
- 13) 宇高健太郎, 早川典子, 北田克己, 森田恒之, 荒井経, 稲葉政満, 半田昌規, 斎藤典彦. 膠の基礎知識. 膠文化研究会; 2015
- 14) UDAKA Kentaro, HAYAKAWA Noriko, KITADA Katsumi, MORITA Tsuneyuki, ARAI Kei, INABA Masamitsu, HANDA Masaki, SAITO Norihiko, MATSUBARA Michiko. Basic Knowledge about Animal Glue. Nikawa Labs; 2017
- 15) 岡岩太郎, 小笠原具子, 佐藤千晴. 粉状化した白色顔料の剥落止めに用いる膠の検討 ～宇高健太郎氏作製古典的膠と中川胡粉製播州粒膠の使用を例に～. 鳳翔学叢第 13 輯. 平等院; 2017
- 16) 早川典子. 平等院鳳凰堂東面中央扉に使用した剥落止め材料について. 鳳翔学叢第 13 輯. 平等院; 2017
- 17) 宇高健太郎. 膠の性状と装潢における適性. 膠文化研究会第 11 回公開研究会「膠と修理 - 《序の舞》を守る-. 予稿集; 2018
- 18) 宇高健太郎. 文化財修復における膠の適性. 膠文化研究会第 12 回公開研究会「膠千年」予稿集; 2019
- 19) 宇野雪村. 必携 文房古玩事典. 柏書房株式会社; 1983
- 20) 編 東京藝術大学大学院文化財保存学日本画研究室. 図解 日本画用語事典. (株)東京美術; 2007

- 21) 編集委員長 加藤時雄. カーボンブラック便覧〈第三版〉. カーボンブラック協会; 1995
- 22) 宇高健太郎. 煤製造における諸条件と製品の性状の関連. 文化財保存修復学会第35回大会研究発表要旨集; 2014
- 23) 宇高健太郎. 煤及び膠に関する研究. 文化財保存修復学会第37回大会研究発表要旨集; 2015
- 24) 宇高健太郎. 松煙煤に関する研究. 文化財保存修復学会第40回大会研究発表要旨; 2018
- 25) UDAKA Kentaro, Jennifer A. Giaccari. Research on Properties of Soot for Inkstick Productions. Collection of research presentations of the 42nd Annual Meeting of the Japan Society for the Conservation of Cultural Property; 2020
- 26) 宇高健太郎. 膠製造における諸条件と製品の性状の関連. 文化財保存修復学会第33回大会研究発表要旨集; 2011
- 27) 宇高健太郎. 膠製造における諸条件と製品の性状の関連(2) -製造条件と界面性状等-. 文化財保存修復学会第34回大会研究発表要旨集; 2012
- 28) 宇高健太郎, 野村義宏, 寺岡葉子. 膠製造における諸条件と製品の性状の関連(3) -製造条件と物性決定因子-. 文化財保存修復学会第34回大会研究発表要旨集; 2012
- 29) 宇高健太郎. 膠製造における諸条件と製品の性状の関連(4). 文化財保存修復学会第35回大会研究発表要旨集; 2013
- 30) 宇高健太郎. 膠製造における諸条件と製品の性状の関連(5). 文化財保存修復学会第36回大会研究発表要旨集; 2014
- 31) 宇高健太郎. 膠の性状に関する研究. 文化財保存修復学会第38回大会研究発表要旨集; 2016
- 32) 宇高健太郎, 早川典子, 半田昌規, 岡泰央, 藤井佑果, 小笠原具子, 亀井亮子, 半田幾子, 宇和川史彦, 柏谷明美. 膠の性状と装潢における適性の関連. 文化財保存修復学会第39回大会研究発表要旨集; 2017
- 33) 宇高健太郎, 早川典子, 藤井佑果, 大場詩野子, 岡部迪子, 柏谷明美. 膠の調製等に関する研究. 文化財保存修復学会第41回大会研究発表要旨集; 2019
- 34) 宇高健太郎, 早川典子, 藤井佑果, 柏谷明美. 膠の性状と湿熱劣化処理の影響に関する研究 -表面観察による検証-. 保存科学 58; 2019
- 35) 宇高健太郎, 早川典子, 藤井佑果, 大場詩野子, 岡部迪子. 古典的膠の調製方法及び性状. 文化財保存修復学会第42回大会研究発表集; 2020
- 36) 鈴木梅仙. 梅仙墨談; 1897
- 37) 松尾良樹、的場美帆. 『唐人墨製問答之記録』 - 「古梅園製墨資料」翻刻と解題(2) -. 古代学 4; 2012
- 38) 濱田薫. 残笏居 古墨記言; 2007
- 39) 訳編 赤堀昭. 中国製墨資料. 呉竹精昇堂; 1986
- 40) 輯 陶湘. 涉園墨萃. 中國書店; 1991 (原典 中華民國期-宋代)
- 41) 渡邊明義. 水墨画の鑑賞基礎知識. (株)至文堂; 1997
- 42) 仙石正. 新・墨談. 内田老鶴園新社; 1971
- 43) 綿谷正之. 墨の文化史 概説. 奈良保育学院研究紀要 16; 2014

- 44) All Japan Handmade Washi Association. Handbook on the Art of Washi. Wagami-do K.K.; 1991
- 45) 編 馬淵久夫, 杉下龍一郎, 三輪嘉六, 沢田正昭, 三浦定俊. 文化財科学の事典. (株)朝倉書店; 2003
- 46) 潘吉星, 訳 佐藤武敏. 中國製紙技術史. (株)平凡社; 1990
- 47) 松井和泉掾元泰, 松井和泉掾元彙, 訳註編 松尾良樹. 古梅園墨譜. (株)古梅園; 1993 (原典 1716-73)
- 48) 編 奈良文化財研究所, 監修 佐藤昌憲. 絹文化財の世界 伝統文化技術と保存科学. 角川学芸出版; 2005
- 49) 森田早織. 青蓮院所蔵国宝「不動明王二童子像」の表現技法に関する研究 -造形表現と技法材料の関連性について-. 東京藝術大学博士研究論文; 2014
- 50) 編集委員長 河合武司. 改訂第2版 界面と界面活性剤 -基礎から応用まで-. (社)日本油化学会; 2009
- 51) 日中美術展実行委員会. 日中国交正常化40周年記念 日中美術展 東洋美術の未来を探る-日本画と工筆画 (於 (株)東京美術倶楽部) 図録; 2012
- 52) SOSU International Co., Ltd. MIHARAYASUHIRO "MAY-DAY", "BOTAN", 2014-15AW Paris Collection; 2014
- 53) 鈴木桂一郎. 紀州生まつ松煙. 新仁商店; 1993
- 54) 松井茂雄. The 墨 -墨は生きている-. (株)日貿出版社; 1983
- 55) 松井藤次. 和墨の伝流考. (株)墨運堂; 1981
- 56) 宋応星, 訳 藪内清. 天工開物. (株)平凡社; 1969 (原典 1637)
- 57) 松井和泉掾元泰, 編 松井貞太郎. 古梅園墨談. (株)古梅園; 1934 (原典 1713)
- 58) 榊莫山. 文房四宝 墨の話. (株)角川学芸出版; 1998
- 59) 植村和堂. 和硯と和墨. 理工学社; 1980
- 60) (株)墨運堂企画室. 墨のQ&A. (株)墨運堂; 2001
- 61) Velson Horie. Materials for Conservation: Organic Consolidants, Adhesives and Coatings, Second edition. Routledge Taylor & Francis Group; 2010
- 62) 編集代表 我孫子義弘. 改訂版にかわとゼラチン-産業史と科学技術-. 日本にかわ・ゼラチン工業組合; 1997
- 63) 森田恒之. 画材の博物誌. 中央公論美術出版; 1994
- 64) 山内章, 木下雅代. 文化財保存修復に用いる和膠の生産研究と今後の展望. 文化財保存修復学会第32回大会研究発表要旨集; 2010
- 65) 青木芳昭. 絵画素材の科学 よくわかる今の絵画材料. (株)生活の友社; 2011
- 66) 杉田正見. 白なめし革の化学的・組織学的解明による製造方法の改善に関する研究; 1986年
- 67) 出口公長. 印伝革の技術史的考察と製造技術に関する研究; 2000
- 68) 荒井経. 日本画と材料 近代に作られた伝統. (株)武蔵野美術大学出版局; 2015
- 69) 古田亮. 日本画とは何だったのか 近代日本画史論. (株)KADOKAWA; 2018
- 70) 推訳註 (有)金開堂, 原著 林守篤. 画筌. (有)金開堂; 2010 (原典 1712)
- 71) 染谷香理. 日本画画材関連史料翻刻集 I (江戸中期篇); 2018
- 72) 推訳註 (有)金開堂, 原著 土佐光起. 本朝画法大伝. (有)金開堂; 2002 (原典 1690)

- 73) 林久良. 日本皮革工芸史研究 正倉院漆皮袈裟箱の復元; 2003
- 74) 訳 西山武一, 熊代幸雄. 校訂譯註 齊民要術. アジア經濟出版会; 1969 (原典 後魏代)
- 75) 森由雄. 神農本草經解説. (有)源草社; 2011 (原典 1854 および後漢代-三国期)
- 76) 原著 テオフィルス, 訳編 森洋. さまざまの技能について. (株)中央公論美術出版; 1996 (原典 1100 頃)

■Chapter 19. Collagen Based Foods

SUZUKI Satoshi

Nippi, Incorporated, Gelatin-related Division

- 1) 和田正汎, 長谷川忠男, コラーゲンとゼラチンの科学, 2011 年
- 2) 安孫子義弘, にかわとゼラチン, 1997 年
- 3) 早出武, コラーゲンケーシング, 食肉の科学, 1996 年; 37:189-194
- 4) 渡部睦人, 野村義宏, コラーゲン今昔物語~可食性フィルム(第6回目), 皮革科学, 2014 年; 60:1-5
- 5) 高井恒昌, お菓子の国からこんにちは, ムース・ババロア・プディング, 1986 年
- 6) 辻薦, 食品加工ハンドブック, 1971 年:239-243
- 7) 鈴木哲, ゼラチンの基本物性と応用アプリケーション, フードケミカル, 2011 年;11:39-44
- 8) 金谷節子, ベッドサイドから在宅で使える 嚥下食のすべて 医歯薬出版株式会社
- 9) 小原仁, 土居守, 摂食・嚥下障害を有する慢性期リハビリテーション患者におけるゼラチンを活用した嚥下障害食の栄養状態に対する効果、栄養学会誌 Vol. 64、No. 4、237-242 (2006 年)
- 10) Adam M. Welche Wirkung haben Gelatinepräparate, Therapiewoche 1991; 38: 2456-61
- 11) Adam M, Spacek P, Hulejova H, Galianova A, Blahos J, Postmenopausal osteoporosis. Treatment with calcitonin and a diet rich in collagen proteins, Cas Lek Cesk, 1996; 135: 74-8
- 12) 菊池数晃, 又平芳春, ヒトの乾燥肌および肌荒れに対する海洋性コラーゲンペプチド含有飲料の有用性, FRAGRANCE JOURNAL, 2004; 2003-9 : 97-102.
- 13) 角田愛美, 広田亜里沙, 桑葉くみ子, 楠畑雅, 小山洋一, 新谷隆行 et al, 皮膚角層の吸水能と血液性化学検査に対するコラーゲンペプチド経口摂取の作用, Journal of Nutritional Food, 2004 年; 7: 45-52
- 14) Aaron M. Altschul, タンパク質 —その化学と栄養政策— 昭和 42 年、光琳書院
- 15) 日本食品標準成分表 2015 年版 (七訂)

■Chapter 20. Nutraceuticals

KOIZUMI Seiko, MATSUSHITA Aya, DAIMON Mona, HIMENO Ai

Research and Development Center, Nitta Gelatin Inc.

- 1) DEPARTMENT OF HEALTH AND HUMAN SERVICES, Food and Drug Administration, 21 CFR Chapter I [Docket No. 02N-0434] Federal Register. 2003; 68(77).
- 2) Husøy T, Bruzell E, Granum B, Hetland RB, Rohloff J, Wicklund T, et al. Risk Assessment of "Other Substances" -Collagen from Fish Skin; *European Journal of Nutrition & Food Safety* 2018; 8(3): 105-107.
- 3) Ikoma T, Kobayashi H, Tanaka J, Walsh D and Mann S, Physical properties of type I collagen extracted from fish scales of *Pagrus major* and *Oreochromis niloticus*; *Int J Biol Macromol.* 2003; 32(3-5): 199-204.
- 4) 梶原苗美, コラーゲン由来ペプチドの栄養効果に関する研究; 神戸女子大学家政学部紀要, 1999; 32: 41-50.
- 5) 小柳達男, 鷹嘴テル, 及川桂子, ゼラチンおよびアミノ酸の補足による米の栄養価改善; *食糧と栄養*, 1966;19(4); 256-259.
- 6) Prockop DJ, Keisher HR, Sjoerdsma A. Gastrointestinal absorption and renal excretion of hydroxyproline peptides; *Lancet.* 1962; 2: 527
- 7) Adam M, Welche Wirkung haben Gelatinepräparate? *Therapiewoche.* 1991; 38: 2456-61.
- 8) Liu C, Sugita K, Nihei K, Yoneyama K, Tanaka H. Absorption of hydroxyproline-containing peptides in vascularly perfused rat small intestine in situ. *Biosci Biotechnol Biochem.* 2009; 73(8): 1741-7.
- 9) Osawa Y, Mizushige T, Jinno S, Sugihara F, Inoue N, Tanaka H. Absorption and metabolism of orally administered collagen hydrolysates evaluated by the vascularly perfused rat intestine and liver in situ. *Biomedical Research (Tokyo)* 2018; 39 (1): 1-11
- 10) Yamamoto S, Deguchi K, Onuma M, Numata N, Sakai Y. Absorption and Urinary Excretion of Peptides after Collagen Tripeptide Ingestion in Humans. *Biol. Pharm. Bull.* 2016; 39, 428-434
- 11) Ichikawa S, Morifuji M, Ohara H, Matsumoto H, Takeuchi Y, Sato K. Hydroxyproline-containing dipeptides and tripeptides quantified at high concentration in human blood after oral administration of gelatin hydrolysate. *Int J Food Sci Nutr.* 2010; 61(1): 52-60.
- 12) Sugihara F, Inoue N, Kuwamori M, Taniguchi M. Quantification of hydroxyprolyl-glycine (Hyp-Gly) in human blood after ingestion of collagen hydrolysate. *J Biosci Bioeng.* 2012; 113(2): 202-3.
- 13) Iwai K, Hasegawa T, Taguchi Y, Morimatsu F, Sato K, Nakamura Y. Identification of food-derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. *J Agric Food Chem.* 2005 Aug 10;53(16): 6531-6.

- 14) Shigemura Y, Kubomura D, Sato Y, Sato K. Dose-dependent changes in the levels of free and peptide forms of hydroxyproline in human plasma after collagen hydrolysate ingestion. *Food Chem.* 2014; 15: 159; 328–32.
- 15) Shigemura Y, Akaba S, Kawashima E, Park EY, Nakamura Y, Sato K. Identification of a novel food-derived collagen peptide, hydroxyprolyl-glycine, in human peripheral blood by pre-column derivatisation with phenyl isothiocyanate. *Food Chem.* 2011; 129(3): 1019–24.
- 16) Kawaguchi T, Nanbu PN, Kurokawa M. Distribution of prolylhydroxyproline and its metabolites after oral administration in rats. *Biol. Pharm. Bull.* 2012; 35: 422–427
- 17) Meilman E, Urivetzky M, and Rapoport M. URINARY HYDROXYPROLINE PEPTIDES; *J Clin Invest.* 1963; 42(1): 40–50.
- 18) Franchesca D, Choi BS R, Calvin T, Sung BS, Margit L.W. Juhasz MD. Oral Collagen Supplementation; A Systematic Review of Dermatological Applications. *J Drugs Dermatol.* 2019; 18(1): 9–16.
- 19) Sibilla S, Godfrey M, Brewer S, Budh-Raja A, Genovese L. An Overview of the Beneficial Effects of Hydrolysed Collagen as a Nutraceutical on Skin Properties; Scientific Background and Clinical Studies. *The Open Nutraceuticals Journal.* 2015; 8: 29–42
- 20) Proksch E, Segger D, Degwert J, Schunck M, Zague V, Oesser S. Oral supplementation of specific collagen peptides has beneficial effects on human skin physiology; a double-blind, placebo-controlled study. *Skin Pharmacol Physiol.* 2014; 27(1): 47–55.
- 21) Sugihara F, Inoue N, Wang X. Clinical effects of ingesting collagen hydrolysate on facial skin properties –A randomized, placebo-controlled, double-blind trial–. *Jpn Pharmacol Ther* 2015; 43(1): 67–70
- 22) Tanaka M, Koyama Y, Nomura Y. Effects of collagen peptide ingestion on UV-B-induced skin damage. *Biosci Biotechnol Biochem.* 2009;73(4); 930–2.
- 23) Ohara H, Ichikawa S, Matsumoto H, Akiyama M, Fujimoto N, Kobayashi T. Collagen-derived dipeptide, proline-hydroxyproline, stimulates cell proliferation and hyaluronic acid synthesis in cultured human dermal fibroblasts. *J. Dermatol.* 2010; 37: 330–338.
- 24) Sugihara F, Inoue N, Venkateswarathirukumara S. Ingestion of bioactive collagen hydrolysates enhanced pressure ulcer healing in a randomized double-blind placebo-controlled clinical study *Scientific Reports.* 2018; 8: 11403
- 25) Koyama Y. Effects of Collagen Ingestion and their Biological Significance; *J Nutr Food Sci.* 2016; 6: 3
- 26) Elam M, Johnson S, Hooshmand S, Feresin R, Payton M, Gu J. A calcium-collagen chelate dietary supplement attenuates bone loss in postmenopausal women with osteopenia; a randomized controlled trial. *J Med Food.* 2015; 18(3): 324–31.
- 27) Koyama Y, Hirota A, Mori H, Takahara H, Kuwaba K, Kusubata M, et al. Ingestion of Gelatin Has

- Differential Effect on Bone Mineral Density and Body Weight in Protein Undernutrition. *Journal of Nutritional Science and Vitaminology*. 2001; 47(1): 84–6.
- 28) Kimira Y, Ogura K, Taniuchi Y, Kataoka A, Inoue N, Sugihara F, et al. Collagen-derived dipeptide prolyl-hydroxyproline promotes differentiation of MC3T3-E1 osteoblastic cells. *Biochemical and Biophysical Research Communications*. 2014; 453(3): 498–501.
- 29) Kimira Y, Odaira H, Nomura K, Taniuchi Y, Inoue N, Nakatani S, et al. Collagen-derived dipeptide prolyl-hydroxyproline promotes osteogenic differentiation through Foxg1. *Cellular & Molecular Biology Letters*. 2017; 22: 27
- 30) Nomura K, Kimira Y, Osawa Y, Shimizu J, Matsushita A, Mano H. Collagen-derived dipeptide prolyl hydroxyproline directly binds to Foxg1 to change its conformation and inhibit the interaction with Runx2. *Biosci Biotechnol Biochem*. 2019; 83(11): 2027–2033.
- 31) 川口浩, 変形性関節症治療の国内外のガイドライン; *日関病誌*; 2016; 35(1); 1–9
- 32) Moskowitz R. Role of collagen hydrolysate in bone and joint disease; *Semin Arthritis Rheum*. 2000; 30(2): 87–99.
- 33) Shen J, Show Y. Collagen peptides improve knee osteoarthritis in elderly women; A 6-month randomized, double-blind, placebo-controlled study. *Agro. Food Industry Hi Tech* 2014; 25(2): 19–23
- 34) Beynen A, Van Geene H, Grim H, Jacobs P, Van T. Oral Administration of Gelatin Hydrolysate Reduces Clinical Signs of Canine Osteoarthritis in a Double-Blind, Placebo-Controlled Trial. *American Journal of Animal and Veterinary Sciences*. 2010; 5(2): 102–6
- 35) McAlindon T, Nuite M, Krishnan N, Ruthazer R, Price L, Burstein D, et al. Change in knee osteoarthritis cartilage detected by delayed gadolinium enhanced magnetic resonance imaging following treatment with collagen hydrolysate; a pilot randomized controlled trial. *Osteoarthritis and Cartilage*. 2011; 19: 399e405
- 36) Kumar S, Sugihara F, Suzuki K, Inoue N, Venkateswarathirukumara S. A double-blind, placebo-controlled, randomised, clinical study on the effectiveness of collagen peptide on osteoarthritis. *J Sci Food Agric*. 2015; 95(4): 702–7.
- 37) Kimira Y, Saito M, Kushibe S, Nomura K, Osawa Y, Sekiguchi Y, et al. The Effects of Collagen Peptide Supplementation on Knee Joint Health—A Double-blind, Placebo-controlled, Randomized Trial in Healthy University Students Belonging to a Running Club—. *Jpn Pharmacol Ther*. 2019; 47(9): 1455 - 1462
- 38) Isaka S, Someya A, Nakamura S, Naito K, Nozawa M, Inoue N, et al. Evaluation of the effect of oral administration of collagen peptides on an experimental rat osteoarthritis model. *Exp Ther Med*. 2017;13(6): 2699–2706.
- 39) Nakatani S, Mano H, Sampei C, Shimizu J, Wada M. Chondroprotective effect of the bioactive peptide prolyl-hydroxyproline in mouse articular cartilage in vitro and in vivo. *Osteoarthritis*

Cartilage. 2009;17(12): 1620–7.

- 40) Ohara H, Iida H, Ito K, Takeuchi Y and Nomura Y. Effects of Pro-Hyp, a collagen hydrolysate-derived peptide, on hyaluronic acid synthesis using in vitro cultured synovium cells and oral ingestion of collagen hydrolysates in a guinea pig model of osteoarthritis. *Biosci Biotechnol Biochem*; 2010; 74(10): 2096–9.
- 41) Zdzieblik D, Oesser S, Baumstark M, Gollhofer A, König D. Collagen peptide supplementation in combination with resistance training improves body composition and increases muscle strength in elderly sarcopenic men; a randomised controlled trial. *Br J Nutr*. 2015; 114(8): 1237–1245.
- 42) Clifford T, Ventress M, Allerton DM, Stansfield S, Tang JCY, Fraser WD, et al. The effects of collagen peptides on muscle damage, inflammation and bone turnover following exercise; a randomized, controlled trial. *Amino Acids*. 2019; 51(4): 691–704
- 43) Kitakaze T, Sakamoto T, Kitano T, Inoue N, Sugihara F, Harada N. The collagen derived dipeptide hydroxyprolyl-glycine promotes C2C12 myoblast differentiation and myotube hypertrophy. *Biochem Biophys Res Commun*. 2016; 478(3): 1292–7.
- 44) Lassoued I, Nasri L, Mourad J, Nasri M. Angiotensin Converting Enzyme Inhibition with Antiradicalar Activity of Thornback Ray Gelatin Hydrolysate; Determination and Separation of Peptide Fractions. *Journal of Hypertension and Management*. 2017; 3(1): 1–8.
- 45) Torkova A, Kononikhin A, Bugrova A, Khotchenkov V, Tsentalovich M, Medvedeva U. Effect of in Vitro Gastrointestinal Digestion on Bioactivity of Poultry Protein Hydrolysate. *Current Research in Nutrition and Food Science* 2016; 4(2): 77–86
- 46) Saiga A, Iwai K, Hayakawa T, Takahata Y, Kitamura S, Nishimura T, et al. Angiotensin I-converting enzyme-inhibitory peptides obtained from chicken collagen hydrolysate. *J Agric Food Chem*. 2008; 56: 9586–9591.
- 47) Tomosugi N, Yamamoto S, Takeuchi M, Yonekura H, Ishigaki Y, Numata N, et al. Effect of Collagen Tripeptide on Atherosclerosis in Healthy Humans. *J Atheroscler Thromb*. 2017; 1; 24(5): 530–538.
- 48) Igase M, Kohara K, Okada Y, Ochi M, Igase K, Inoue N, et al. A double-blind, placebo-controlled, randomised clinical study of the effect of pork collagen peptide supplementation on atherosclerosis in healthy older individuals. *Biosci Biotechnol Biochem*. 2018; 15:1–3.
- 49) Kouguchi T, Ohmori T, Shimizu M, Takahata Y, Maeyama Y, Suzuki T, et al. Effects of a chicken collagen hydrolysate on the circulation system in subjects with mild hypertension or high-normal blood pressure. *Biosci Biotechnol Biochem*. 2013;77(4): 691–696.
- 50) Devasia S, Kumar S, Stephen PS, Inoue N, Sugihara F, Suzuki K. Double Blind, Randomized Clinical Study to Evaluate Efficacy of Collagen Peptide as Add on Nutritional Supplement in Type 2 Diabetes. *Clin Nutr Food Sci*. 2018; 1(1): 006–11.
- 51) Devasia S, Kumar S, PS S, Inoue N, Sugihara F, Koizumi S, et al, A Double Blind, Randomised,

Four Arm Clinical Study to Evaluate the Safety, Efficacy and Tolerability of Collagen Peptide as a Nutraceutical Therapy in the Management of Type II Diabetes Mellitus; *Journal of Diabetes & Metabolism*; 2019; 10(11), No. 839; 1-7.

- 52) Koizumi S, Inoue N, Sugihara F, Igase M, Effects of Collagen Hydrolysates on Human Brain Structure and Cognitive Function: A Pilot Clinical Study. *Nutrients*. 2019; 12(1); E50.
- 53) Nishikimi A, Koyama Y, Ishihara S, Kobayashi S, Tometsuka C, Kusubata M, et al. Collagen-derived peptides modulate CD4+ T-cell differentiation and suppress allergic responses in mice. *Immun Inflamm Dis*. 2018; 6(2): 245-255.

■Chapter 21. Collagen in the Cosmetics Field

AMANO Satoshi

Shiseido Global Innovation Center

- 1) Ohara H, Ichikawa S, Matsumoto H, et al. Collagen-derived dipeptide, proline-hydroxyproline, stimulates cell proliferation and hyaluronic acid synthesis in cultured human dermal fibroblasts. *J Dermatol* 2010;37:330-8.
- 2) Birk DE, Fitch JM, Babiarz JP, Doane KJ, Linsenmayer TF. Collagen fibrillogenesis in vitro: interaction of types I and V collagen regulates fibril diameter. *J Cell Sci* 1990;95 (Pt 4):649-57.
- 3) Keene DR, Sakai LY, Bachinger HP, Burgeson RE. Type III collagen can be present on banded collagen fibrils regardless of fibril diameter. *J Cell Biol* 1987;105:2393-402.
- 4) Nishiyama T, McDonough AM, Bruns RR, Burgeson RE. Type XII and XIV collagens mediate interactions between banded collagen fibers in vitro and may modulate extracellular matrix deformability. *J Biol Chem* 1994;269:28193-9.
- 5) Theocharidis G, Drymoussi Z, Kao AP, et al. Type VI Collagen Regulates Dermal Matrix Assembly and Fibroblast Motility. *J Invest Dermatol* 2016;136:74-83.
- 6) Amano S, Matsunaga Y, Akutsu N, et al. Basement membrane damage, a sign of skin early aging, and laminin 5, a key player in basement membrane care. *IFSCC Magazine* 2000;3:15-23.
- 7) Fisher GJ, Datta SC, Talwar HS, et al. Molecular basis of sun-induced premature skin ageing and retinoid antagonism. *Nature* 1996;379:335-9.
- 8) Takada K, Amano S, Kohno Y, Nishiyama T, Inomata S. Non-invasive study of gelatinases in sun-exposed and unexposed healthy human skin based on measurements in stratum corneum. *Arch Dermatol Res* 2006;298:237-42.
- 9) Inomata S, Matsunaga Y, Amano S, et al. Possible involvement of gelatinases in basement membrane damage and wrinkle formation in chronically ultraviolet B-exposed hairless mouse. *J Invest Dermatol* 2003;120:128-34.
- 10) Hachiya A, Sriwiriyanont P, Fujimura T, et al. Mechanistic effects of long-term ultraviolet B irradiation induce epidermal and dermal changes in human skin xenografts. *Am J Pathol* 2009;174:401-13.
- 11) Iriyama S, Matsunaga Y, Takahashi K, Matsuzaki K, Kumagai N, Amano S. Activation of heparanase by ultraviolet B irradiation leads to functional loss of basement membrane at the dermal-epidermal junction in human skin. *Arch Dermatol Res* 2011;303:253-61.
- 12) Iriyama S, Hiruma T, Tsunenaga M, Amano S. Influence of heparan sulfate chains in proteoglycan at the dermal-epidermal junction on epidermal homeostasis. *Exp Dermatol* 2011;20:810-4.
- 13) Iriyama S, Matsunaga Y, Amano S. Heparanase activation induces epidermal hyperplasia,

- angiogenesis, lymphangiogenesis and wrinkles. *Exp Dermatol* 2010;19:965–72.
- 14) Tsunenaga M, Kohno Y, Horii I, et al. Growth and differentiation properties of normal and transformed human keratinocytes in organotypic culture. *Jpn J Cancer Res* 1994;85:238–44.
 - 15) Amano S, Akutsu N, Matsunaga Y, et al. Importance of balance between extracellular matrix synthesis and degradation in basement membrane formation. *Exp Cell Res* 2001;271:249–62.
 - 16) Amano S, Nishiyama T, Burgeson RE. A specific and sensitive ELISA for laminin 5. *J Immunol Methods* 1999;224:161–9.
 - 17) Amano S, Ogura Y, Akutsu N, Nishiyama T. Quantitative analysis of the synthesis and secretion of type VII collagen in cultured human dermal fibroblasts with a sensitive sandwich enzyme-linked immunoassay. *Exp Dermatol* 2007;16:151–5.
 - 18) Akutsu N, Amano S, Nishiyama T. Quantitative analysis of laminin 5 gene expression in human keratinocytes. *Exp Dermatol* 2005;14:329–35.
 - 19) Ogura Y, Matsunaga Y, Nishiyama T, Amano S. Plasmin induces degradation and dysfunction of laminin 332 (laminin 5) and impaired assembly of basement membrane at the dermal-epidermal junction. *Br J Dermatol* 2008;159:49–60.
 - 20) Tsunenaga M, Adachi E, Amano S, Burgeson RE, Nishiyama T. Laminin 5 can promote assembly of the lamina densa in the skin equivalent model. *Matrix Biol* 1998;17:603–13.
 - 21) Iriyama S, Tsunenaga M, Amano S, Adachi E. Key role of heparan sulfate chains in assembly of anchoring complex at the dermal-epidermal junction. *Exp Dermatol* 2011;20:953–5.
 - 22) Iriyama S, Yamanishi H, Kunizawa N, Hirao T, Amano S. 1-(2-Hydroxyethyl)-2-imidazolidinone, a heparanase and matrix metalloproteinase inhibitor, improves epidermal basement membrane structure and epidermal barrier function. *Exp Dermatol* 2019;28:247–53.
 - 23) Oba A, Gomi, T., Nishimori, T., Graves, C., Pearse, A., and Edwards, C. Non-invasive methods for assessment of dermal UV damage. *J Soc Cosmet Chem* 2004;38:193–201.
 - 24) Fisher GJ, Varani J, Voorhees JJ. Looking older: fibroblast collapse and therapeutic implications. *Arch Dermatol* 2008;144:666–72.
 - 25) Agarwal P, Zwolanek D, Keene DR, et al. Collagen XII and XIV, new partners of cartilage oligomeric matrix protein in the skin extracellular matrix suprastructure. *J Biol Chem* 2012;287:22549–59.
 - 26) Kobayashi M, Kawabata K, Kusaka-Kikushima A, et al. Cartilage Oligomeric Matrix Protein Increases in Photodamaged Skin. *J Invest Dermatol* 2016;136:1143–9.
 - 27) Kielty CM, Sherratt MJ, Shuttleworth CA. Elastic fibres. *J Cell Sci* 2002;115:2817–28.
 - 28) Kadoya K, Amano S, Nishiyama T, et al. Changes in fibrillin-1 expression, elastin expression and skin surface texture at sites of cultured epithelial autograft transplantation onto wounds from burn scar excision. *International wound journal* 2015.
 - 29) Yano K, Kadoya K, Kajiya K, Hong YK, Detmar M. Ultraviolet B irradiation of human skin induces

an angiogenic switch that is mediated by upregulation of vascular endothelial growth factor and by downregulation of thrombospondin-1. *Br J Dermatol* 2005;152:115-21.

- 30) Ezure T, Amano S. Increment of subcutaneous adipose tissue is associated with decrease of elastic fibres in the dermal layer. *Exp Dermatol* 2015;24:924-9.
- 31) Ezure T, Amano S. Adiponectin and leptin up-regulate extracellular matrix production by dermal fibroblasts. *Biofactors* 2007;31:229-36.
- 32) Ezure T, Amano S. Negative regulation of dermal fibroblasts by enlarged adipocytes through release of free fatty acids. *J Invest Dermatol* 2011;131:2004-9.

■Chapter 22. Medical Collagen

FUJIMOTO Ichiro

Koken Research Institute Co., Ltd.

- 1) Miller EJ, Lane JM, Piez KA. Isolation and characterization of the peptides derived from the alpha-1 chain of chick bone collagen after cyanogen bromide cleavage. *Biochemistry*. 1969 Jan;8(1):30-9.
- 2) Veit G, Kobbe B, Keene DR, Paulsson M, Koch M, Wagener R. Collagen XXVIII, a novel von Willebrand factor A domain-containing protein with many imperfections in the collagenous domain. *J Biol Chem*. 2006 Feb 10;281(6):3494-504.
- 3) https://www.mhlw.go.jp/web/t_doc?dataId=81004000&dataType=0&pageNo=1
- 4) Boyce ST, Kagan RJ, Meyer NA, Yakuboff KP, Warden GD. The 1999 clinical research award. Cultured skin substitutes combined with Integra Artificial Skin to replace native skin autograft and allograft for the closure of excised full-thickness burns. *J Burn Care Rehabil*. 1999 Nov-Dec;20(6):453-61.
- 5) Morimoto N, Yoshimura K, Niimi M, Ito T, Aya R, Fujitaka J, Tada H, Teramukai S, Murayama T, Toyooka C, Miura K, Takemoto S, Kanda N, Kawai K, Yokode M, Shimizu A, Suzuki S. Novel collagen/gelatin scaffold with sustained release of basic fibroblast growth factor: clinical trial for chronic skin ulcers. *Tissue Eng Part A*. 2013 Sep;19(17-18):1931-40.
- 6) Takata T, Wang HL, Miyauchi M. Attachment, proliferation and differentiation of periodontal ligament cells on various guided tissue regeneration membranes. *J Periodontal Res*. 2001 Oct;36(5):322-7.
- 7) Nakamura T, Inada Y, Fukuda S, Yoshitani M, Nakada A, Itoi S, Kanemaru S, Endo K, Shimizu Y. Experimental study on the regeneration of peripheral nerve gaps through a polyglycolic acid-collagen (PGA-collagen) tube. *Brain Res*. 2004 Nov 19;1027(1-2):18-29.
- 8) Sugaya, A., Minabe, M., Hori, T., Tatsumi, J., Watanabe, Y., Ikeda, K., Numabe, Y., Hsyashi, H. and Kamoi, K. : Effects on wound healing of tricalcium phosphate - collagen complex implants in periodontal osseous defects in the dog. *J. Periodontal. Res.*, 1990 Jan; 25(1): 60-3.
- 9) Browder IW, Litwin MS. Use of absorbable collagen for hemostasis in general surgical patients. *Am Surg*. 1986 Sep;52(9):492-4.
- 10) Aebi M. N-linked protein glycosylation in the ER. *Biochim Biophys Acta*. 2013 Nov;1833(11):2430-7.